

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46397 A2(51) International Patent Classification⁷: C12N 9/12

(21) International Application Number: PCT/US00/35304

(22) International Filing Date:
20 December 2000 (20.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/172,066 23 December 1999 (23.12.1999) US
60/176,107 14 January 2000 (14.01.2000) US
60/177,731 21 January 2000 (21.01.2000) US
60/178,573 28 January 2000 (28.01.2000) USSunnyvale, CA 94086 (US). YUE, Henry [US/US]; 826
Lois Avenue, Sunnyvale, CA 94087 (US). YAO, Monique,
G. [US/US]; 111 Frederick Court, Mountain View, CA
94043 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa
Clara, CA 95054 (US). KHAN, Farrah, A. [IN/US]; 333
Escuela Avenue #221, Mountain View, CA 94040 (US).(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics,
Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(71) Applicant (*for all designated States except US*): INCYTE
GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo
Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): YANG, Jun-
ming [CN/US]; 7125 Bark Lane, San Jose, CA 95129
(US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago
Road, San Leandro, CA 94577 (US). BURFORD, Neil
[GB/US]; 105 Wildwood Circle, Durham, CT 06422
(US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle
Lane, Brisbane, CA 94005 (US). LU, Dyung, Aina, M.
[US/US]; 233 Coy Drive, San Jose, CA 95123 (US).
REDDY, Roopa [IN/US]; 1233 W. McKinley Avenue #3,(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished
upon receipt of that report.For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HUMAN KINASES

(57) Abstract: The invention provides human kinases (PKIN) and polynucleotides which identify and encode PKIN. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of PKIN.

HUMAN KINASES

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of human kinases and to the use of these sequences in the diagnosis, treatment, and prevention of cancer, immune disorders, disorders affecting growth and development, cardiovascular diseases, and lipid disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of human kinases.

BACKGROUND OF THE INVENTION

Kinases comprise the largest known enzyme superfamily and vary widely in their target molecules. Kinases catalyze the transfer of high energy phosphate groups from a phosphate donor to a phosphate acceptor. Nucleotides usually serve as the phosphate donor in these reactions, with most kinases utilizing adenosine triphosphate (ATP). The phosphate acceptor can be any of a variety of molecules, including nucleosides, nucleotides, lipids, carbohydrates, and proteins. Proteins are phosphorylated on hydroxyamino acids. Addition of a phosphate group alters the local charge on the acceptor molecule, causing internal conformational changes and potentially influencing intermolecular contacts. Reversible protein phosphorylation is the primary method for regulating protein activity in eukaryotic cells. In general, proteins are activated by phosphorylation in response to extracellular signals such as hormones, neurotransmitters, and growth and differentiation factors. The activated proteins initiate the cell's intracellular response by way of intracellular signaling pathways and second messenger molecules such as cyclic nucleotides, calcium-calmodulin, inositol, and various mitogens, that regulate protein phosphorylation.

Kinases are involved in all aspects of a cell's function, from basic metabolic processes, such as glycolysis, to cell-cycle regulation, differentiation, and communication with the extracellular environment through signal transduction cascades. Inappropriate phosphorylation of proteins in cells has been linked to changes in cell cycle progression and cell differentiation. Changes in the cell cycle have been linked to induction of apoptosis or cancer. Changes in cell differentiation have been linked to diseases and disorders of the reproductive system, immune system, and skeletal muscle.

There are two classes of protein kinases. One class, protein tyrosine kinases (PTKs), phosphorylates tyrosine residues, and the other class, protein serine/threonine kinases (STKs), phosphorylates serine and threonine residues. Some PTKs and STKs possess structural characteristics of both families and have dual specificity for both tyrosine and serine/threonine residues. Almost all kinases contain a conserved 250-300 amino acid catalytic domain containing specific residues and

sequence motifs characteristic of the kinase family. The protein kinase catalytic domain can be further divided into 11 subdomains. N-terminal subdomains I-IV fold into a two-lobed structure which binds and orients the ATP donor molecule, and subdomain V spans the two lobes. C-terminal subdomains VI-XI bind the protein substrate and transfer the gamma phosphate from ATP to the hydroxyl group of a tyrosine, serine, or threonine residue. Each of the 11 subdomains contains specific catalytic residues or amino acid motifs characteristic of that subdomain. For example, subdomain I contains an 8-amino acid glycine-rich ATP binding consensus motif, subdomain II contains a critical lysine residue required for maximal catalytic activity, and subdomains VI through IX comprise the highly conserved catalytic core. PTKs and STKs also contain distinct sequence motifs in subdomains VI and VIII which may confer hydroxyamino acid specificity.

In addition, kinases may also be classified by additional amino acid sequences, generally between 5 and 100 residues, which either flank or occur within the kinase domain. These additional amino acid sequences regulate kinase activity and determine substrate specificity. (Reviewed in Hardie, G. and Hanks, S. (1995) The Protein Kinase Facts Book, Vol I p.p. 17-20 Academic Press, San Diego, CA.). In particular, two protein kinase signature sequences have been identified in the kinase domain, the first containing an active site lysine residue involved in ATP binding, and the second containing an aspartate residue important for catalytic activity. If a protein analyzed includes the two protein kinase signatures, the probability of that protein being a protein kinase is close to 100% (PROSITE: PDOC00100, November 1995).

20 Protein Tyrosine Kinases

Protein tyrosine kinases (PTKs) may be classified as either transmembrane, receptor PTKs or nontransmembrane, nonreceptor PTK proteins. Transmembrane tyrosine kinases function as receptors for most growth factors. Growth factors bind to the receptor tyrosine kinase (RTK), which causes the receptor to phosphorylate itself (autophosphorylation) and specific intracellular second messenger proteins. Growth factors (GF) that associate with receptor PTKs include epidermal GF, platelet-derived GF, fibroblast GF, hepatocyte GF, insulin and insulin-like GFs, nerve GF, vascular endothelial GF, and macrophage colony stimulating factor.

Nontransmembrane, nonreceptor PTKs lack transmembrane regions and, instead, form signaling complexes with the cytosolic domains of plasma membrane receptors. Receptors that function through non-receptor PTKs include those for cytokines and hormones (growth hormone and prolactin), and antigen-specific receptors on T and B lymphocytes.

Many PTKs were first identified as oncogene products in cancer cells in which PTK activation was no longer subject to normal cellular controls. In fact, about one third of the known oncogenes encode PTKs. Furthermore, cellular transformation (oncogenesis) is often accompanied by increased

tyrosine phosphorylation activity (Charbonneau, H. and Tonks, N. K. (1992) *Annu. Rev. Cell Biol.* 8:463-93). Regulation of PTK activity may therefore be an important strategy in controlling some types of cancer.

Protein Serine/Threonine Kinases

5 Protein serine/threonine kinases (STKs) are nontransmembrane proteins. A subclass of STKs are known as ERKs (extracellular signal regulated kinases) or MAPs (mitogen-activated protein kinases) and are activated after cell stimulation by a variety of hormones and growth factors. Cell stimulation induces a signaling cascade leading to phosphorylation of MEK (MAP/ERK kinase) which, in turn, activates ERK via serine and threonine phosphorylation. A varied number of proteins represent
10 the downstream effectors for the active ERK and implicate it in the control of cell proliferation and differentiation, as well as regulation of the cytoskeleton. Activation of ERK is normally transient, and cells possess dual specificity phosphatases that are responsible for its down-regulation. Also, numerous studies have shown that elevated ERK activity is associated with some cancers. Other STKs include the second messenger dependent protein kinases such as the cyclic-AMP dependent protein kinases
15 (PKA), calcium-calmodulin (CaM) dependent protein kinases, and the mitogen-activated protein kinases (MAP); the cyclin-dependent protein kinases; checkpoint and cell cycle kinases; proliferation-related kinases; 5'-AMP-activated protein kinases; and kinases involved in apoptosis.

The second messenger dependent protein kinases primarily mediate the effects of second messengers such as cyclic AMP (cAMP), cyclic GMP, inositol triphosphate, phosphatidylinositol,
20 3,4,5-triphosphate, cyclic ADPribose, arachidonic acid, diacylglycerol and calcium-calmodulin. The PKAs are involved in mediating hormone-induced cellular responses and are activated by cAMP produced within the cell in response to hormone stimulation. cAMP is an intracellular mediator of hormone action in all animal cells that have been studied. Hormone-induced cellular responses include thyroid hormone secretion, cortisol secretion, progesterone secretion, glycogen breakdown,
25 bone resorption, and regulation of heart rate and force of heart muscle contraction. PKA is found in all animal cells and is thought to account for the effects of cAMP in most of these cells. Altered PKA expression is implicated in a variety of disorders and diseases including cancer, thyroid disorders, diabetes, atherosclerosis, and cardiovascular disease (Isselbacher, K.J. et al. (1994) Harrison's Principles of Internal Medicine, McGraw-Hill, New York, NY, pp. 416-431, 1887).

30 The casein kinase I (CKI) gene family is another subfamily of serine/threonine protein kinases. This continuously expanding group of kinases have been implicated in the regulation of numerous cytoplasmic and nuclear processes, including cell metabolism, and DNA replication and repair. CKI enzymes are present in the membranes, nucleus, cytoplasm and cytoskeleton of eukaryotic cells, and on the mitotic spindles of mammalian cells (Fish, K.J. et al., (1995) *J. Biol. Chem.* 270:14875-14883).

The CKI family members all have a short amino-terminal domain of 9-76 amino acids, a highly conserved kinase domain of 284 amino acids, and a variable carboxyl-terminal domain that ranges from 24 to over 200 amino acids in length (Cegielska, A. et al., (1998) J. Biol. Chem. 273:1357-1364.) The CKI family is comprised of highly related proteins, as seen by the identification of isoforms of casein kinase I from a variety of sources. There are at least five mammalian isoforms, α , β , γ , δ , and ϵ . Fish et al., identified CKI-epsilon from a human placenta cDNA library. It is a basic protein of 416 amino acids and is closest to CKI-delta. Through recombinant expression, it was determined to phosphorylate known CKI substrates and was inhibited by the CKI-specific inhibitor CKI-7. The human gene for CKI-epsilon was able to rescue yeast with a slow-growth phenotype caused by deletion of the yeast CKI locus, HRR250 (Fish et al, *supra*.)

The mammalian circadian mutation tau was found to be a semidominant autosomal allele of CKI-epsilon that markedly shortens period length of circadian rhythms in Syrian hamsters. The tau locus is encoded by casein kinase I-epsilon, which is also a homolog of the *Drosophila* circadian gene double-time. Studies of both the wildtype and tau mutant CKI-epsilon enzyme indicated that the mutant enzyme has a noticeable reduction in the maximum velocity and autophosphorylation state. Further, *in vitro*, CKI-epsilon is able to interact with mammalian PERIOD proteins, while the mutant enzyme is deficient in its ability to phosphorylate PERIOD. Lowrey et al., have proposed that CKI-epsilon plays a major role in delaying the negative feedback signal within the transcription-translation-based autoregulatory loop that composes the core of the circadian mechanism. Therefore the CKI-epsilon enzyme is an ideal target for pharmaceutical compounds influencing circadian rhythms, jet-lag and sleep, in addition to other physiologic and metabolic processes under circadian regulation (Lowrey, P.L. et al., (2000) Science 288:483-491.)

Calcium-Calmodulin Dependent Protein Kinases

Calcium-calmodulin dependent (CaM) kinases are involved in regulation of smooth muscle contraction, glycogen breakdown (phosphorylase kinase), and neurotransmission (CaM kinase I and CaM kinase II). CaM dependent protein kinases are activated by calmodulin, an intracellular calcium receptor, in response to the concentration of free calcium in the cell. Many CaM kinases are also activated by phosphorylation. Some CaM kinases are also activated by autophosphorylation or by other regulatory kinases. CaM kinase I phosphorylates a variety of substrates including the neurotransmitter-related proteins synapsin I and II, the gene transcription regulator, CREB, and the cystic fibrosis conductance regulator protein, CFTR (Haribabu, B. et al. (1995) EMBO Journal 14:3679-3686). CaM kinase II also phosphorylates synapsin at different sites and controls the synthesis of catecholamines in the brain through phosphorylation and activation of tyrosine hydroxylase. CaM kinase II controls the synthesis of catecholamines and serotonin, through

phosphorylation/activation of tyrosine hydroxylase and tryptophan hydroxylase, respectively (Fujisawa, H. (1990) *BioEssays* 12:27-29). The mRNA encoding a calmodulin-binding protein kinase-like protein was found to be enriched in mammalian forebrain. This protein is associated with vesicles in both axons and dendrites and accumulates largely postnatally. The amino acid sequence of this protein is similar to

5 CaM-dependent STKs, and the protein binds calmodulin in the presence of calcium (Godbout, M. et al. (1994) *J. Neurosci.* 14:1-13).

Mitogen-Activated Protein Kinases

The mitogen-activated protein kinases (MAP) which mediate signal transduction from the cell surface to the nucleus via phosphorylation cascades are another STK family that regulates intracellular

10 signaling pathways. Several subgroups have been identified, and each manifests different substrate specificities and responds to distinct extracellular stimuli (Egan, S.E. and Weinberg, R.A. (1993) *Nature* 365:781-783). MAP kinase signaling pathways are present in mammalian cells as well as in yeast. The extracellular stimuli which activate MAP kinase pathways include epidermal growth factor (EGF), ultraviolet light, hyperosmolar medium, heat shock, endotoxic lipopolysaccharide (LPS), and

15 pro-inflammatory cytokines such as tumor necrosis factor (TNF) and interleukin-1 (IL-1). Altered MAP kinase expression is implicated in a variety of disease conditions including cancer, inflammation, immune disorders, and disorders affecting growth and development.

Cyclin-Dependent Protein Kinases

The cyclin-dependent protein kinases (CDKs) are STKs that control the progression of cells

20 through the cell cycle. The entry and exit of a cell from mitosis are regulated by the synthesis and destruction of a family of activating proteins called cyclins. Cyclins are small regulatory proteins that bind to and activate CDKs, which then phosphorylate and activate selected proteins involved in the mitotic process. CDKs are unique in that they require multiple inputs to become activated. In addition to cyclin binding, CDK activation requires the phosphorylation of a specific threonine residue and the

25 dephosphorylation of a specific tyrosine residue on the CDK.

Another family of STKs associated with the cell cycle are the NIMA (never in mitosis)-related kinases (Neks). Both CDKs and Neks are involved in duplication, maturation, and separation of the microtubule organizing center, the centrosome, in animal cells (Fry, A.M., et al. (1998) *EMBO J.* 17:470-481).

Checkpoint and Cell Cycle Kinases

In the process of cell division, the order and timing of cell cycle transitions are under control of cell cycle checkpoints, which ensure that critical events such as DNA replication and chromosome segregation are carried out with precision. If DNA is damaged, e.g. by radiation, a checkpoint pathway is activated that arrests the cell cycle to provide time for repair. If the damage is extensive, apoptosis is

induced. In the absence of such checkpoints, the damaged DNA is inherited by aberrant cells which may cause proliferative disorders such as cancer. Protein kinases play an important role in this process. For example, a specific kinase, checkpoint kinase 1 (Chk1), has been identified in yeast and mammals, and is activated by DNA damage in yeast. Activation of Chk1 leads to the arrest of the cell at the G2/M transition. (Sanchez, Y. et al. (1997) Science 277:1497-1501.) Specifically, Chk1 phosphorylates the cell division cycle phosphatase CDC25, inhibiting its normal function which is to dephosphorylate and activate the cyclin-dependent kinase Cdc2. Cdc2 activation controls the entry of cells into mitosis. (Peng, C-Y et al. (1997) Science 277:1501- 1505.) Thus, activation of Chk1 prevents the damaged cell from entering mitosis. A similar deficiency in a checkpoint kinase, such as Chk1, may also contribute to cancer by failure to arrest cells with damaged DNA at other checkpoints such as G2/M.

Proliferation-Related Kinases

Proliferation-related kinase is a serum/cytokine inducible STK that is involved in regulation of the cell cycle and cell proliferation in human megakaryocytic cells (Li, B. et al. (1996) J. Biol. Chem. 271:19402-8). Proliferation-related kinase is related to the polo (derived from *Drosophila* polo gene) family of STKs implicated in cell division. Proliferation-related kinase is downregulated in lung tumor tissue and may be a proto-oncogene whose deregulated expression in normal tissue leads to oncogenic transformation.

5'-AMP-activated protein kinase

A ligand-activated STK protein kinase is 5'-AMP-activated protein kinase (AMPK) (Gao, G. et al. (1996) J. Biol. Chem. 271:8675-8681). Mammalian AMPK is a regulator of fatty acid and sterol synthesis through phosphorylation of the enzymes acetyl-CoA carboxylase and hydroxymethylglutaryl-CoA reductase and mediates responses of these pathways to cellular stresses such as heat shock and depletion of glucose and ATP. AMPK is a heterotrimeric complex comprised of a catalytic alpha subunit and two non-catalytic beta and gamma subunits that are believed to regulate the activity of the alpha subunit. Subunits of AMPK have a much wider distribution in non-lipogenic tissues such as brain, heart, spleen, and lung than expected. This distribution suggests that its role may extend beyond regulation of lipid metabolism alone.

Kinases in Apoptosis

Apoptosis is a highly regulated signaling pathway leading to cell death that plays a crucial role in tissue development and homeostasis. Deregulation of this process is associated with the pathogenesis of a number of diseases including autoimmune disease, neurodegenerative disorders, and cancer. Various STKs play key roles in this process. ZIP kinase is an STK containing a C-terminal leucine zipper domain in addition to its N-terminal protein kinase domain. This C-terminal domain appears to

mediate homodimerization and activation of the kinase as well as interactions with transcription factors such as activating transcription factor, ATF4, a member of the cyclic-AMP responsive element binding protein (ATF/CREB) family of transcriptional factors (Sanjo, H. et al. (1998) J. Biol. Chem. 273:29066-29071). DRAK1 and DRAK2 are STKs that share homology with the death-associated protein kinases (DAP kinases), known to function in interferon- γ induced apoptosis (Sanjo et al. supra). Like ZIP kinase, DAP kinases contain a C-terminal protein-protein interaction domain, in the form of ankyrin repeats, in addition to the N-terminal kinase domain. ZIP, DAP, and DRAK kinases induce morphological changes associated with apoptosis when transfected into NIH3T3 cells (Sanjo et al. supra). However, deletion of either the N-terminal kinase catalytic domain or the C-terminal domain of these proteins abolishes apoptosis activity, indicating that in addition to the kinase activity, activity in the C-terminal domain is also necessary for apoptosis, possibly as an interacting domain with a regulator or a specific substrate.

RICK is another STK recently identified as mediating a specific apoptotic pathway involving the death receptor, CD95 (Inohara, N. et al. (1998) J. Biol. Chem. 273:12296-12300). CD95 is a member of the tumor necrosis factor receptor superfamily and plays a critical role in the regulation and homeostasis of the immune system (Nagata, S. (1997) Cell 88:355-365). The CD95 receptor signaling pathway involves recruitment of various intracellular molecules to a receptor complex following ligand binding. This process includes recruitment of the cysteine protease caspase-8 which, in turn, activates a caspase cascade leading to cell death. RICK is composed of an N-terminal kinase catalytic domain and a C-terminal "caspase-recruitment" domain that interacts with caspase-like domains, indicating that RICK plays a role in the recruitment of caspase-8. This interpretation is supported by the fact that the expression of RICK in human 293T cells promotes activation of caspase-8 and potentiates the induction of apoptosis by various proteins involved in the CD95 apoptosis pathway (Inohara et al. supra).

Mitochondrial Protein Kinases

A novel class of eukaryotic kinases, related by sequence to prokaryotic histidine protein kinases, are the mitochondrial protein kinases (MPKs) which seem to have no sequence similarity with other eukaryotic protein kinases. These protein kinases are located exclusively in the mitochondrial matrix space and may have evolved from genes originally present in respiration-dependent bacteria which were endocytosed by primitive eukaryotic cells. MPKs are responsible for phosphorylation and inactivation of the branched-chain alpha-ketoacid dehydrogenase and pyruvate dehydrogenase complexes (Harris, R.A. et al. (1995) Adv. Enzyme Regul. 34:147-162). Five MPKs have been identified. Four members correspond to pyruvate dehydrogenase kinase isozymes, regulating the activity of the pyruvate dehydrogenase complex, which is an important regulatory enzyme at the interface between glycolysis and the citric acid cycle. The fifth member corresponds to a branched-

chain alpha-ketoacid dehydrogenase kinase, important in the regulation of the pathway for the disposal of branched-chain amino acids. (Harris, R.A. et al. (1997) Adv. Enzyme Regul. 37:271-293). Both starvation and the diabetic state are known to result in a great increase in the activity of the pyruvate dehydrogenase kinase in the liver, heart and muscle of the rat. This increase contributes in both disease states to the phosphorylation and inactivation of the pyruvate dehydrogenase complex and conservation of pyruvate and lactate for gluconeogenesis (Harris (1995) *supra*).

KINASES WITH NON-PROTEIN SUBSTRATES

10 Lipid and Inositol kinases

Lipid kinases phosphorylate hydroxyl residues on lipid head groups. A family of kinases involved in phosphorylation of phosphatidylinositol (PI) has been described, each member phosphorylating a specific carbon on the inositol ring (Leever, S.J. et al. (1999) Curr. Opin. Cell. Biol. 11:219-225). The phosphorylation of phosphatidylinositol is involved in activation of the protein kinase C signaling pathway. The inositol phospholipids (phosphoinositides) intracellular signaling pathway begins with binding of a signaling molecule to a G-protein linked receptor in the plasma membrane. This leads to the phosphorylation of phosphatidylinositol (PI) residues on the inner side of the plasma membrane by inositol kinases, thus converting PI residues to the biphosphate state (PIP₂). PIP₂ is then cleaved into inositol triphosphate (IP₃) and diacylglycerol. These two products act as mediators for separate signaling pathways. Cellular responses that are mediated by these pathways are glycogen breakdown in the liver in response to vasopressin, smooth muscle contraction in response to acetylcholine, and thrombin-induced platelet aggregation.

PI 3-kinase (PI3K), which phosphorylates the D3 position of PI and its derivatives, has a central role in growth factor signal cascades involved in cell growth, differentiation, and metabolism. PI3K is a heterodimer consisting of an adapter subunit and a catalytic subunit. The adapter subunit acts as a scaffolding protein, interacting with specific tyrosine-phosphorylated proteins, lipid moieties, and other cytosolic factors. When the adapter subunit binds tyrosine phosphorylated targets, such as the insulin responsive substrate (IRS)-1, the catalytic subunit is activated and converts PI (4,5) biphosphate (PIP₂) to PI (3,4,5) P₃ (PIP₃). PIP₃ then activates a number of other proteins, including PKA, protein kinase B (PKB), protein kinase C (PKC), glycogen synthase kinase (GSK)-3, and p70 ribosomal s6 kinase. PI3K also interacts directly with the cytoskeletal organizing proteins, Rac, rho, and cdc42 (Shepherd, P.R., et al. (1998) Biochem. J. 333:471-490). Animal models for diabetes, such as *obese* and *fat* mice, have altered PI3K adapter subunit levels. Specific mutations in the adapter subunit have also been found in an insulin-resistant Danish population, suggesting a role for PI3K in

type-2 diabetes (Shepard, supra).

An example of lipid kinase phosphorylation activity is the phosphorylation of D-erythro-sphingosine to the sphingolipid metabolite, sphingosine-1-phosphate (SPP). SPP has emerged as a novel lipid second-messenger with both extracellular and intracellular actions (Kohama, T. et al. (1998) J. Biol. Chem. 273:23722-23728). Extracellularly, SPP is a ligand for the G-protein coupled receptor EDG-1 (endothelial-derived, G-protein coupled receptor). Intracellularly, SPP regulates cell growth, survival, motility, and cytoskeletal changes. SPP levels are regulated by sphingosine kinases that specifically phosphorylate D-erythro-sphingosine to SPP. The importance of sphingosine kinase in cell signaling is indicated by the fact that various stimuli, including platelet-derived growth factor (PDGF), nerve growth factor, and activation of protein kinase C, increase cellular levels of SPP by activation of sphingosine kinase, and the fact that competitive inhibitors of the enzyme selectively inhibit cell proliferation induced by PDGF (Kohama et al. supra).

Purine Nucleotide Kinases

The purine nucleotide kinases, adenylate kinase (ATP:AMP phosphotransferase, or AdK) and guanylate kinase (ATP:GMP phosphotransferase, or GuK) play a key role in nucleotide metabolism and are crucial to the synthesis and regulation of cellular levels of ATP and GTP, respectively. These two molecules are precursors in DNA and RNA synthesis in growing cells and provide the primary source of biochemical energy in cells (ATP), and signal transduction pathways (GTP). Inhibition of various steps in the synthesis of these two molecules has been the basis of many antiproliferative drugs for cancer and antiviral therapy (Pillwein, K. et al. (1990) Cancer Res. 50:1576-1579).

AdK is found in almost all cell types and is especially abundant in cells having high rates of ATP synthesis and utilization such as skeletal muscle. In these cells AdK is physically associated with mitochondria and myofibrils, the subcellular structures that are involved in energy production and utilization, respectively. Recent studies have demonstrated a major function for AdK in transferring high energy phosphoryls from metabolic processes generating ATP to cellular components consuming ATP (Zelevnikar, R.J. et al. (1995) J. Biol. Chem. 270:7311-7319). Thus AdK may have a pivotal role in maintaining energy production in cells, particularly those having a high rate of growth or metabolism such as cancer cells, and may provide a target for suppression of its activity to treat certain cancers. Alternatively, reduced AdK activity may be a source of various metabolic, muscle-energy disorders that can result in cardiac or respiratory failure and may be treatable by increasing AdK activity.

GuK, in addition to providing a key step in the synthesis of GTP for RNA and DNA synthesis, also fulfills an essential function in signal transduction pathways of cells through the regulation of GDP and GTP. Specifically, GTP binding to membrane associated G proteins mediates the activation of cell

- receptors, subsequent intracellular activation of adenylyl cyclase, and production of the second messenger, cyclic AMP. GDP binding to G proteins inhibits these processes. GDP and GTP levels also control the activity of certain oncogenic proteins such as p21^{ras} known to be involved in control of cell proliferation and oncogenesis (Bos, J.L. (1989) Cancer Res. 49:4682-4689). High ratios of
- 5 GTP:GDP caused by suppression of GuK cause activation of p21^{ras} and promote oncogenesis. Increasing GuK activity to increase levels of GDP and reduce the GTP:GDP ratio may provide a therapeutic strategy to reverse oncogenesis.

- GuK is an important enzyme in the phosphorylation and activation of certain antiviral drugs useful in the treatment of herpes virus infections. These drugs include the guanine homologs acyclovir and bucidovir (Miller, W.H. and Miller R.L. (1980) J. Biol. Chem. 255:7204-7207; Stenberg, K. et al. (1986) J. Biol. Chem. 261:2134-2139). Increasing GuK activity in infected cells may provide a therapeutic strategy for augmenting the effectiveness of these drugs and possibly for reducing the necessary dosages of the drugs.

Pyrimidine Kinases

- 15 The pyrimidine kinases are deoxycytidine kinase and thymidine kinase 1 and 2. Deoxycytidine kinase is located in the nucleus, and thymidine kinase 1 and 2 are found in the cytosol (Johansson, M. et al. (1997) Proc. Natl. Acad. Sci. U.S.A. 94:11941-11945). Phosphorylation of deoxyribonucleosides by pyrimidine kinases provides an alternative pathway for de novo synthesis of DNA precursors. The role of pyrimidine kinases, like purine kinases, in phosphorylation is critical to the activation of several
- 20 chemotherapeutically important nucleoside analogues (Arner E.S. and Eriksson, S. (1995) Pharmacol. Ther. 67:155-186).

- The discovery of new human kinases and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cancer, immune disorders, disorders affecting growth and development, cardiovascular diseases, and
- 25 lipid disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of human kinases.

SUMMARY OF THE INVENTION

- The invention features purified polypeptides, human kinases, referred to collectively as "PKIN"
- 30 and individually as "PKIN-1," "PKIN-2," "PKIN-3," "PKIN-4," "PKIN-5," "PKIN-6," "PKIN-7," "PKIN-8," "PKIN-9," "PKIN-10," "PKIN-11," and "PKIN-12." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the

group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-12.

- 5 The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of
- 10 SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-12. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:13-24.

- Additionally, the invention provides a recombinant polynucleotide comprising a promoter
- 15 sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID
- 20 NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

- The invention also provides a method for producing a polypeptide comprising an amino acid
- 25 sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group
- 30 consisting of SEQ ID NO:1-12. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

 Additionally, the invention provides an isolated antibody which specifically binds to a

polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or

absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence
5 selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and a pharmaceutically acceptable excipient. In one
10 embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional PKIN, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an
15 agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino
20 acid sequence selected from the group consisting of SEQ ID NO:1-12. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of
25 functional PKIN, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence
30 selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising

an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional PKIN, comprising administering to a patient in need of such treatment the composition.

5 The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence selected
10 from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

15 The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, c) a biologically active fragment of an amino acid sequence
20 selected from the group consisting of SEQ ID NO:1-12, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with
25 the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

 The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a
30 sequence selected from the group consisting of SEQ ID NO:13-24, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

 The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound;

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, ii) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, ii) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

20

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for each polypeptide of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of each polypeptide sequence, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of each polypeptide.

Table 4 lists the cDNA and genomic DNA fragments which were used to assemble each polynucleotide sequence, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for each polynucleotide of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

5 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

10 It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

15 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, 20 protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"PKIN" refers to the amino acid sequences of substantially purified PKIN obtained from any 25 species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of PKIN. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PKIN either by directly interacting with 30 PKIN or by acting on components of the biological pathway in which PKIN participates.

An "allelic variant" is an alternative form of the gene encoding PKIN. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to

allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding PKIN include those sequences with deletions, 5 insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as PKIN or a polypeptide with at least one functional characteristic of PKIN. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding PKIN, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding PKIN. The 10 encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent PKIN. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of PKIN is retained. For example, negatively charged amino acids may 15 include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

20 The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to a sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

25 “Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of PKIN. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small 30 molecules, or any other compound or composition which modulates the activity of PKIN either by directly interacting with PKIN or by acting on components of the biological pathway in which PKIN participates.

The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant.

Antibodies that bind PKIN polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers
5 that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which
10 bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA;
15 peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once
20 introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical
25 functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic PKIN, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement,
30 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding PKIN or fragments of PKIN may be

employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

5 "Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap
10 (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino
15 acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
	Arg	His, Lys
20	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
25	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
30	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
35	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation,
40 (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of PKIN or the polynucleotide encoding PKIN which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:13-24 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:13-24, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:13-24 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:13-24 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:13-24 and the region of SEQ ID NO:13-24 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-12 is encoded by a fragment of SEQ ID NO:13-24. A fragment of SEQ ID NO:1-12 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-12. For example, a fragment of SEQ ID NO:1-12 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-12. The precise length of a fragment of SEQ ID NO:1-12 and the region of SEQ ID NO:1-12 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended

purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

5 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in
10 the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular
15 biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between
20 aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at
25 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The
30 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

5 *Expect: 10*

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over
10 the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

15 Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to
20 the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

25 Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with
30 polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

5 *Word Size: 3*

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

15 "Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

20 "Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic

strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of PKIN which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of PKIN which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of PKIN. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of PKIN.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an PKIN may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of PKIN.

"Probe" refers to nucleic acid sequences encoding PKIN, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100,

or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence

that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have
5 been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is
10 expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

15 "Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear
20 sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nucleogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing PKIN, nucleic acids encoding PKIN, or fragments thereof may comprise a bodily fluid; an extract from a cell,
25 chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure
30 of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are

removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

5 A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

10 A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences
15 into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed
20 cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor
25 of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be
30 introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at

- least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule.
- 10 Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be
- 15 indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

- A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence
- 20 identity over a certain defined length of one of the polypeptides.

THE INVENTION

- The invention is based on the discovery of new human kinases (PKIN), the polynucleotides encoding PKIN, and the use of these compositions for the diagnosis, treatment, or prevention of cancer,
- 25 immune disorders, disorders affecting growth and development, cardiovascular diseases, and lipid disorders.

- Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.
- 30

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of each of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:13-24 or that distinguish between SEQ ID NO:13-24 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 2287966H1 is the identification number of an Incyte cDNA sequence, and BRAINON01 is the cDNA library from which

it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 70166939V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g2821547) which contributed to the assembly of the full length polynucleotide sequences. Alternatively, the identification numbers in column 5 may refer to coding regions predicted by Genscan analysis of genomic DNA. For example, g4454511.v113.gs_3.nt.edit is the identification number of a Genscan-predicted coding sequence, with g4454511 being the GenBank identification number of the sequence to which Genscan was applied. The Genscan-predicted coding sequences may have been edited prior to assembly. (See Example IV.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. (See Example V.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon-stretching" algorithm. (See Example V.) In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses PKIN variants. A preferred PKIN variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the PKIN amino acid sequence, and which contains at least one functional or structural characteristic of PKIN.

The invention also encompasses polynucleotides which encode PKIN. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:13-24, which encodes PKIN. The polynucleotide sequences of SEQ ID NO:13-24, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding PKIN. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding PKIN. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:13-24 which has at

least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:13-24. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of PKIN.

5 It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding PKIN, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in
10 accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring PKIN, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode PKIN and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring PKIN under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding PKIN or its
15 derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PKIN and its derivatives without altering the encoded amino acid sequences include the production of RNA
20 transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode PKIN and PKIN derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using
25 reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding PKIN or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:13-24 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and
30 S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of

DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding PKIN may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been

size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

5 Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate
10 software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

 In another embodiment of the invention, polynucleotide sequences or fragments thereof which
15 encode PKIN may be cloned in recombinant DNA molecules that direct expression of PKIN, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express PKIN.

 The nucleotide sequences of the present invention can be engineered using methods generally
20 known in the art in order to alter PKIN-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites,
25 alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

 The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or
30 improve the biological properties of PKIN, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of

DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of
5 homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding PKIN may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively,
10 PKIN itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp.55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be
15 achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of PKIN, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid
20 chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active PKIN, the nucleotide sequences encoding PKIN or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains
25 the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding PKIN. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding PKIN. Such signals
30 include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding PKIN and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the

vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

- 5 Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding PKIN and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

- 10 A variety of expression vector/host systems may be utilized to contain and express sequences encoding PKIN. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus);
- 15 plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO*
- 20 *J.* 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di
- 25 Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.)
- The invention is not limited by the host cell employed.

- 30 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding PKIN. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding PKIN can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding PKIN into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed

bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of PKIN are needed, e.g. for the production of antibodies,
5 vectors which direct high level expression of PKIN may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of PKIN. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such
10 vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of PKIN. Transcription of sequences encoding
15 PKIN may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can
20 be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding PKIN may be ligated into an
25 adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses PKIN in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-
30 based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of PKIN in cell lines is preferred. For example, sequences encoding PKIN can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding PKIN is inserted within a marker gene sequence, transformed cells containing sequences encoding PKIN can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PKIN under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding PKIN and that express PKIN may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and

protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of PKIN using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PKIN is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding PKIN include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding PKIN, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding PKIN may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode PKIN may be designed to contain signal sequences which direct secretion of PKIN through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells

which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

5 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding PKIN may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric PKIN protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of PKIN activity. Heterologous protein and peptide moieties
10 may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins,
15 respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the PKIN encoding sequence and the heterologous protein sequence, so that PKIN may be cleaved away from the heterologous moiety following purification. Methods for fusion protein
20 expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled PKIN may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or
25 SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

PKIN of the present invention or fragments thereof may be used to screen for compounds that specifically bind to PKIN. At least one and up to a plurality of test compounds may be screened for specific binding to PKIN. Examples of test compounds include antibodies, oligonucleotides, proteins
30 (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of PKIN, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which PKIN

binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express PKIN, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing PKIN or cell membrane fractions which contain PKIN are then contacted with a test compound and binding, stimulation, or inhibition of activity of either PKIN or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with PKIN, either in solution or affixed to a solid support, and detecting the binding of PKIN to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

PKIN of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of PKIN. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for PKIN activity, wherein PKIN is combined with at least one test compound, and the activity of PKIN in the presence of a test compound is compared with the activity of PKIN in the absence of the test compound. A change in the activity of PKIN in the presence of the test compound is indicative of a compound that modulates the activity of PKIN. Alternatively, a test compound is combined with an in vitro or cell-free system comprising PKIN under conditions suitable for PKIN activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of PKIN may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding PKIN or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by

homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding PKIN may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding PKIN can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding PKIN is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress PKIN, e.g., by secreting PKIN in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of PKIN and human kinases. In addition, the expression of PKIN is closely associated with cancers, cell proliferation and cardiovascular diseases. Therefore, PKIN appears to play a role in cancer, immune disorders, disorders affecting growth and development, cardiovascular diseases, and lipid disorders. In the treatment of disorders associated with increased PKIN expression or activity, it is desirable to decrease the expression or activity of PKIN. In the treatment of disorders associated with decreased PKIN expression or activity, it is desirable to increase the expression or activity of PKIN.

Therefore, in one embodiment, PKIN or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PKIN. Examples of such disorders include, but are not limited to, a cancer, such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal

- tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus, leukemias such as multiple myeloma and lymphomas such as Hodgkin's disease; an immune disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis,
- 5 anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis,
- 10 hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial,
- 15 fungal, parasitic, protozoal, and helminthic infections, and trauma; a growth and developmental disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of
- 20 the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus, renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and
- 25 mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss; a cardiovascular disease, such as arteriovenous fistula, atherosclerosis,
- 30 hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular

- calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation, congenital lung anomalies, atelectasis, pulmonary
- 5 congestion and edema, pulmonary embolism, pulmonary hemorrhage, pulmonary infarction, pulmonary hypertension, vascular sclerosis, obstructive pulmonary disease, restrictive pulmonary disease, chronic obstructive pulmonary disease, emphysema, chronic bronchitis, bronchial asthma, bronchiectasis, bacterial pneumonia, viral and mycoplasmal pneumonia, lung abscess, pulmonary tuberculosis, diffuse interstitial diseases, pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative
- 10 interstitial pneumonitis, hypersensitivity pneumonitis, pulmonary eosinophilia bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, Goodpasture's syndromes, idiopathic pulmonary hemosiderosis, pulmonary involvement in collagen-vascular disorders, pulmonary alveolar proteinosis, lung tumors, inflammatory and noninflammatory pleural effusions, pneumothorax, pleural tumors, drug-induced lung disease, radiation-induced lung disease, and complications of lung
- 15 transplantation; and a lipid disorder such as fatty liver, cholestasis, primary biliary cirrhosis, carnitine deficiency, carnitine palmitoyltransferase deficiency, myoadenylate deaminase deficiency, hypertriglyceridemia, lipid storage disorders such as Fabry's disease, Gaucher's disease, Niemann-Pick's disease, metachromatic leukodystrophy, adrenoleukodystrophy, GM₂ gangliosidosis, and ceroid lipofuscinosis, abetalipoproteinemia, Tangier disease, hyperlipoproteinemia, diabetes mellitus,
- 20 lipodystrophy, lipomatosis, acute panniculitis, disseminated fat necrosis, adiposis dolorosa, lipoid adrenal hyperplasia, minimal change disease, lipomas, atherosclerosis, hypercholesterolemia, hypercholesterolemia with hypertriglyceridemia, primary hypoalphalipoproteinemia, hypothyroidism, renal disease, liver disease, lecithin:cholesterol acyltransferase deficiency, cerebrotendinous xanthomatosis, sitosterolemia, hypocholesterolemia, Tay-Sachs disease, Sandhoff's disease,
- 25 hyperlipidemia, hyperlipemia, lipid myopathies, and obesity.

In another embodiment, a vector capable of expressing PKIN or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PKIN including, but not limited to, those described above.

- In a further embodiment, a composition comprising a substantially purified PKIN in
- 30 conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PKIN including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of PKIN may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity

of PKIN including, but not limited to, those listed above.

In a further embodiment, an antagonist of PKIN may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PKIN. Examples of such disorders include, but are not limited to, those cancers, immune disorders, disorders affecting growth and development, cardiovascular diseases, and lipid disorders described above. In one aspect, an antibody which specifically binds PKIN may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express PKIN.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding PKIN may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PKIN including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of PKIN may be produced using methods which are generally known in the art. In particular, purified PKIN may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind PKIN. Antibodies to PKIN may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with PKIN or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PKIN have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at

least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of PKIN amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

- 5 Monoclonal antibodies to PKIN may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and
10 Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

- In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda,
15 S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce PKIN-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

- 20 Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

- Antibody fragments which contain specific binding sites for PKIN may also be generated. For
25 example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

- 30 Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between PKIN and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two

non-interfering PKIN epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for PKIN. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of PKIN-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogenous in their affinities for multiple PKIN epitopes, represents the average affinity, or avidity, of the antibodies for PKIN. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular PKIN epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the PKIN-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of PKIN, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of PKIN-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding PKIN, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding PKIN. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PKIN. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g.,

Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other
 5 gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding PKIN may be used for
 10 somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475),
 15 cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated
 20 cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the
 25 case where a genetic deficiency in PKIN expression or regulation causes disease, the expression of PKIN from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in PKIN are treated by constructing mammalian expression vectors encoding PKIN and introducing these vectors
 30 by mechanical means into PKIN-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J.-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of PKIN include, but are not limited to, the pCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). PKIN may be expressed

5 using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the

10 ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. *supra*), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding PKIN from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID

15 TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these

20 standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to PKIN expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding PKIN under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive

25 element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for

30 receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a

method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference.

Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding PKIN to cells which have one or more genetic abnormalities with respect to the expression of PKIN. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu. Rev. Nutr.* 19:511-544 and Verma, I.M. and N. Somia (1997) *Nature* 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding PKIN to target cells which have one or more genetic abnormalities with respect to the expression of PKIN. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing PKIN to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al. (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary

skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding PKIN to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for PKIN into the alphavirus genome in place of the capsid-coding region results in the production of a large number of PKIN-coding RNAs and the synthesis of high levels of PKIN in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of PKIN into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding PKIN.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by

scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding PKIN. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding PKIN. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased PKIN expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding PKIN may be therapeutically useful, and in the treatment of disorders associated with decreased PKIN expression or activity, a compound which specifically promotes expression of the polynucleotide encoding PKIN may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method

commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding PKIN is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding PKIN are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding PKIN. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various

formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of PKIN, antibodies to PKIN, and mimetics, agonists, antagonists, or inhibitors of PKIN.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising PKIN or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, PKIN or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example PKIN or fragments thereof, antibodies of PKIN, and agonists, antagonists or inhibitors of PKIN, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by

calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind PKIN may be used for the diagnosis of disorders characterized by expression of PKIN, or in assays to monitor patients being treated with PKIN or agonists, antagonists, or inhibitors of PKIN. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for PKIN include methods which utilize the antibody and a label to detect PKIN in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring PKIN, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PKIN expression. Normal or standard values for PKIN expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to PKIN under

conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of PKIN expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

- 5 In another embodiment of the invention, the polynucleotides encoding PKIN may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of PKIN may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of
- 10 PKIN, and to monitor regulation of PKIN levels during therapeutic intervention.

- In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding PKIN or closely related molecules may be used to identify nucleic acid sequences which encode PKIN. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a
- 15 conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding PKIN, allelic variants, or related sequences.

- Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the PKIN encoding sequences. The hybridization probes of the subject
- 20 invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:13-24 or from genomic sequences including promoters, enhancers, and introns of the PKIN gene.

- Means for producing specific hybridization probes for DNAs encoding PKIN include the cloning of polynucleotide sequences encoding PKIN or PKIN derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to
- 25 synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

- Polynucleotide sequences encoding PKIN may be used for the diagnosis of disorders associated
- 30 with expression of PKIN. Examples of such disorders include, but are not limited to, a cancer, such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus, leukemias such as multiple

- myeloma and lymphomas such as Hodgkin's disease; an immune disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy
- 5 (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis,
- 10 osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a growth and developmental disorder, such as actinic keratosis,
- 15 arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver,
- 20 lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus, renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary
- 25 keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss; a cardiovascular disease, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and
- 30 phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial

- thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation, congenital lung anomalies, atelectasis, pulmonary congestion and edema, pulmonary embolism, pulmonary hemorrhage, pulmonary infarction, pulmonary
- 5 hypertension, vascular sclerosis, obstructive pulmonary disease, restrictive pulmonary disease, chronic obstructive pulmonary disease, emphysema, chronic bronchitis, bronchial asthma, bronchiectasis, bacterial pneumonia, viral and mycoplasmal pneumonia, lung abscess, pulmonary tuberculosis, diffuse interstitial diseases, pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis, pulmonary eosinophilia bronchiolitis
- 10 obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, Goodpasture's syndromes, idiopathic pulmonary hemosiderosis, pulmonary involvement in collagen-vascular disorders, pulmonary alveolar proteinosis, lung tumors, inflammatory and noninflammatory pleural effusions, pneumothorax, pleural tumors, drug-induced lung disease, radiation-induced lung disease, and complications of lung transplantation; and a lipid disorder such as fatty liver, cholestasis, primary biliary cirrhosis, carnitine
- 15 deficiency, carnitine palmitoyltransferase deficiency, myoadenylate deaminase deficiency, hypertriglyceridemia, lipid storage disorders such as Fabry's disease, Gaucher's disease, Niemann-Pick's disease, metachromatic leukodystrophy, adrenoleukodystrophy, GM₂ gangliosidosis, and ceroid lipofuscinosis, abetalipoproteinemia, Tangier disease, hyperlipoproteinemia, diabetes mellitus, lipodystrophy, lipomatosis, acute panniculitis, disseminated fat necrosis, adiposis dolorosa, lipoid
- 20 adrenal hyperplasia, minimal change disease, lipomas, atherosclerosis, hypercholesterolemia, hypercholesterolemia with hypertriglyceridemia, primary hypoalphalipoproteinemia, hypothyroidism, renal disease, liver disease, lecithin:cholesterol acyltransferase deficiency, cerebrotendinous xanthomatosis, sitosterolemia, hypocholesterolemia, Tay-Sachs disease, Sandhoff's disease, hyperlipidemia, hyperlipemia, lipid myopathies, and obesity. The polynucleotide sequences encoding
- 25 PKIN may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered PKIN expression. Such qualitative or quantitative methods are well known in the art.

- In a particular aspect, the nucleotide sequences encoding PKIN may be useful in assays that
- 30 detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding PKIN may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control

sample then the presence of altered levels of nucleotide sequences encoding PKIN in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

- 5 In order to provide a basis for the diagnosis of a disorder associated with expression of PKIN, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PKIN, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with
- 10 values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

- Once the presence of a disorder is established and a treatment protocol is initiated,
- 15 hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

- With respect to cancer, the presence of an abnormal amount of transcript (either under- or
- 20 overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

- 25 Additional diagnostic uses for oligonucleotides designed from the sequences encoding PKIN may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding PKIN, or a fragment of a polynucleotide complementary to the polynucleotide encoding PKIN, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may
- 30 also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

 In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding PKIN may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease

in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding PKIN are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of PKIN include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, PKIN, fragments of PKIN, or antibodies specific for PKIN may be

used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by
5 quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of
10 transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies,
15 or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental
20 compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties.
25 These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data
30 after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is

important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for PKIN to quantify the levels of PKIN expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendozc, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of

methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation
5 between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

10 In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound
15 in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized
20 by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci.*
25 USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

30 In another embodiment of the invention, nucleic acid sequences encoding PKIN may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal

mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

10 Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding PKIN on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA

15 associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is

20 valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may

25 also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

 In another embodiment of the invention, PKIN, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a

30 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between PKIN and the agent being tested may be measured.

 Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are

synthesized on a solid substrate. The test compounds are reacted with PKIN, or fragments thereof, and washed. Bound PKIN is then detected by methods well known in the art. Purified PKIN can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

5 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PKIN specifically compete with a test compound for binding PKIN. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PKIN.

10 In additional embodiments, the nucleotide sequences which encode PKIN may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, 15 therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, in particular U.S. Ser. No. 60/172,066, U.S. Ser. No. 60/176,107, U.S. Ser. No. 60/177,731, and U.S. Ser. No. 60/178,573, are expressly incorporated by reference herein.

20

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. The Incyte cDNA shown for SEQ 25 ID NO:13 was derived from a cDNA library constructed from musculoskeletal tissue. The Incyte cDNA shown for SEQ ID NO:14 was derived from cDNA libraries constructed from prostate, brain and ovarian tissues, including tissues associated with brain, prostate and thyroid tumors. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic 30 solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated

using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

- 5 In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic
- 10 oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,
- 15 PBLUESCRIPT plasmid (Stratagene), PSORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent E. coli cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

20 II. Isolation of cDNA Clones

- Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL
- 25 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4 °C.

- Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal
- 30 cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

- Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the
- 5 MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI
- 10 PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.
- 15 The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS,
- 20 DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank
- 25 cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length
- 30 polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using

MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent

5 identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:13-24. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative human kinases were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode human kinases, the encoded polypeptides were analyzed by querying against PFAM models for kinases. Potential human kinases were also identified by homology to Incyte cDNA sequences that had been annotated as kinases. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information

was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding

5 sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped
10 to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in
15 the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the
20 longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpr public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit
25 from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

"Stretched" Sequences

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public
30 databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the

chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to
5 determine whether it contained a complete gene.

VI. Chromosomal Mapping of PKIN Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:13-24 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched
10 SEQ ID NO:13-24 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment
15 of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

Map locations are represented by ranges, or intervals, or human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in
20 humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified
25 disease genes map within or in proximity to the intervals indicated above.

VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel (1995)
30 supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar.

The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

5

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding PKIN are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding PKIN. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

VIII. Extension of PKIN Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate

fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing

media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

- The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

15 IX. Labeling and Use of Individual Hybridization Probes

- Hybridization probes derived from SEQ ID NO:13-24 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

- The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing

photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra.), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schna (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schna, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated

using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

5 Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia
10 Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and
15 coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic
20 apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in
25 0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with
30 an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide
5 containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477,
10 Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

15 The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different
20 fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC
25 computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

30 A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

XI. Complementary Polynucleotides

Sequences complementary to the PKIN-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring PKIN. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with
5 smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of PKIN. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the PKIN-encoding transcript.

10 XII. Expression of PKIN

Expression and purification of PKIN is achieved using bacterial or virus-based expression systems. For expression of PKIN in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the
15 T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express PKIN upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of PKIN in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as
20 baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding PKIN by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional
25 genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, PKIN is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton
30 enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from PKIN at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-

His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified PKIN obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII where applicable.

5 XIII. Functional Assays

PKIN function is assessed by expressing the sequences encoding PKIN at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which
10 contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the
15 recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include
20 changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated
25 Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of PKIN on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding PKIN and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human
30 immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding PKIN and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIV. Production of PKIN Specific Antibodies

PKIN substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

- 5 Alternatively, the PKIN amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)
- 10 Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide
- 15 and anti-PKIN activity by, for example, binding the peptide or PKIN to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XV. Purification of Naturally Occurring PKIN Using Specific Antibodies

Naturally occurring or recombinant PKIN is substantially purified by immunoaffinity chromatography using antibodies specific for PKIN. An immunoaffinity column is constructed by

20 covalently coupling anti-PKIN antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

- Media containing PKIN are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PKIN (e.g., high ionic strength buffers in the
- 25 presence of detergent). The column is eluted under conditions that disrupt antibody/PKIN binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and PKIN is collected.

XVI. Identification of Molecules Which Interact with PKIN

- PKIN, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent.
- 30 (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled PKIN, washed, and any wells with labeled PKIN complex are assayed. Data obtained using different concentrations of PKIN are used to calculate values for the number, affinity, and association of PKIN with the candidate molecules.

Alternatively, molecules interacting with PKIN are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989) Nature 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

- PKIN may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT)
- 5 which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

XVII. Demonstration of PKIN Activity

- Generally, protein kinase activity is measured by quantifying the phosphorylation of a protein
- 10 substrate by PKIN in the presence of gamma-labeled ^{32}P -ATP. PKIN is incubated with the protein substrate, ^{32}P -ATP, and an appropriate kinase buffer. The ^{32}P incorporated into the substrate is separated from free ^{32}P -ATP by electrophoresis and the incorporated ^{32}P is counted using a radioisotope counter. The amount of incorporated ^{32}P is proportional to the activity of PKIN. A determination of the specific amino acid residue phosphorylated is made by phosphoamino acid analysis of the
- 15 hydrolyzed protein.

- In one alternative, protein kinase activity is measured by quantifying the transfer of gamma phosphate from adenosine triphosphate (ATP) to a serine, threonine or tyrosine residue in a protein substrate. The reaction occurs between a protein kinase sample with a biotinylated peptide substrate and gamma ^{32}P -ATP. Following the reaction, free avidin in solution is added for binding to the
- 20 biotinylated ^{32}P -peptide product. The binding sample then undergoes a centrifugal ultrafiltration process with a membrane which will retain the product-avidin complex and allow passage of free gamma ^{32}P -ATP. The reservoir of the centrifuged unit containing the ^{32}P -peptide product as retentate is then counted in a scintillation counter. This procedure allows assay of any type of protein kinase sample, depending on the peptide substrate and kinase reaction buffer selected. This assay is provided
- 25 in kit form (ASUA, Affinity Ultrafiltration Separation Assay, Transbio Corporation, Baltimore MD, U.S. Patent No. 5,869,275). Suggested substrates and their respective enzymes are as follows: Histone H1 (Sigma) and p34^{cdc2} kinase, Annexin I, Angiotensin (Sigma) and EGF receptor kinase, Annexin II and src kinase, ERK1 & ERK2 substrates and MEK, and myelin basic protein and ERK (Pearson, J.D. et al. (1991) Methods in Enzymology 200:62-81).

- 30 In another alternative, protein kinase activity of PKIN is demonstrated *in vitro* in an assay containing PKIN, 50 μl of kinase buffer, 1 μg substrate, such as myelin basic protein (MBP) or synthetic peptide substrates, 1 mM DTT, 10 μg ATP, and 0.5 μCi [γ - ^{32}P]ATP. The reaction is incubated at 30°C for 30 minutes and stopped by pipetting onto P81 paper. The unincorporated [γ - ^{32}P]ATP is removed by washing and the incorporated radioactivity is measured using a radioactivity scintillation

counter. Alternatively, the reaction is stopped by heating to 100°C in the presence of SDS loading buffer and visualized on a 12% SDS polyacrylamide gel by autoradiography. Incorporated radioactivity is corrected for reactions carried out in the absence of PKIN or in the presence of the inactive kinase, K38A.

- 5 In yet another alternative, adenylate kinase or guanylate kinase activity may be measured by the incorporation of ³²P from gamma-labeled ³²P -ATP into ADP or GDP using a gamma radioisotope counter. The enzyme, in a kinase buffer, is incubated together with the appropriate nucleotide mono-phosphate substrate (AMP or GMP) and ³²P-labeled ATP as the phosphate donor. The reaction is incubated at 37°C and terminated by addition of trichloroacetic acid. The acid extract is neutralized
10 and subjected to gel electrophoresis to separate the mono-, di-, and triphosphonucleotide fractions. The diphosphonucleotide fraction is cut out and counted. The radioactivity recovered is proportional to the enzyme activity.

- In yet another alternative, other assays for PKIN include scintillation proximity assays (SPA), scintillation plate technology and filter binding assays. Useful substrates include recombinant proteins
15 tagged with glutathione transferase, or synthetic peptide substrates tagged with biotin. Inhibitors of PKIN activity, such as small organic molecules, proteins or peptides, may be identified by such assays.

XVIII. Enhancement/Inhibition of Protein Kinase Activity

- Agonists or antagonists of PKIN activation or inhibition may be tested using assays described in section XVII. Agonists cause an increase in PKIN activity and antagonists cause a decrease in PKIN
20 activity.

- Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be
25 understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
058860	1	058860CD1	13	058860CB1
2041716	2	2041716CD1	14	2041716CB1
7472005	3	7472005CD1	15	7472005CB1
7472006	4	7472006CD1	16	7472006CB1
2902460	5	2902460CD1	17	2902460CB1
6383934	6	6383934CD1	18	6383934CB1
3210906	7	3210906CD1	19	3210906CB1
3339024	8	3339024CD1	20	3339024CB1
4436929	9	4436929CD1	21	4436929CB1
5046791	10	5046791CD1	22	5046791CB1
1416174	11	1416174CD1	23	1416174CB1
3244919	12	3244919CD1	24	3244919CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability Score	GenBank Homolog
1	058860CD1	g2677788	8.6e-50	Unknown [Sparisoma chrysotermum], related to g4322024, myosin light chain kinase isoform 3B
2	2041716CD1	g1836161	8.3e-253	Ca ²⁺ /calmodulin-dependent protein kinase IV kinase isoform [Rattus sp.]
3	7472005CD1	g1750259	0.0	Eph-and Elk-related kinase [Mus musculus]
4	7472006CD1	g404634	3.6e-163	Serine/threonine kinase [Mus musculus]
5	2902460CD1	g396429	4.9e-264	IP3 3-kinase [Rattus norvegicus]
6	6383934CD1	g2738898	5.2e-173	Protein kinase [Mus musculus]
7	3210906CD1	g5616074	0.0	Prostate derived STE20-like kinase PSK [Homo sapiens]
8	3339024CD1	g5295850	4.4e-123	QA79 membrane protein [Homo sapiens] (Falco, M. et al. (1999) J. Exp. Med. 190:793-802)
9	4436929CD1	g1872546	0.0	NIK (Nck Interacting Kinase) [Mus musculus] (Su, Y.C. et al. (1997) EMBO J. 16:1279-1290)
10	5046791CD1	g861314	2.7e-21	Similar to Ser/Thr protein kinase [Caenorhabditis elegans]
11	1416174CD1	g8248287	2.00E-61	sphingosine kinase type 2 isoform [Mus musculus]
12	3244919CD1	g7161864	3.10E-185	serine/threonine protein kinase [Mus musculus]

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	058860CD1	466	T422 T5 T12 S19 T31 S46 S83 S168 S179 T194 T331 S351 S365 T422 T52 S163 T299 T312 S402 T451 Y446	N59 N81 N361 N452	Receptor tyrosine kinase: F395-G418 Thiol protease His motif: M116-A126	MOTIFS BLIMPS- BLOCKS
2	2041716CD1	513	S74 T108 S466 T26 S74 S82 S117 S427 S433 T438 T58 S69 S100 S169 S338 S445	N156	ATP/GTP-binding site motif A (P-loop): G493-S500 Serine/Threonine protein kinase active-site signature: I279-L291 Eukaryotic protein kinase domain: Q145-V417 Tyrosine kinase catalytic domain: Y273-L291, G320-I330, L342-D364 Kinase protein beta: M1-Q127 Protein kinase domain: L130-V408	MOTIFS BLAST-DOMO HMMER-PFAM BLIMPS- PRINTS BLAST- PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	7472005CD1	1012	S56 T104 T117 S129 S136 T155 T219 S225 S374 S577 T615 T805 S817 T843 S856 S857 S897 S926 T941 S177 S196 T242 T489 T494 T531 T674 S848 S908 S948 T997 Y487 Y610 Y756	N340 N407 N432 N718 N841	Eukaryotic protein kinase domain: I635-V896 Protein kinases ATP-binding region signature: I641-K667 Tyrosine protein kinases specific active-site signature: Y756-V768 Receptor tyrosine kinase class V: C247-E267 (signature 2) E31-H52, D61-P112, K165-V218, P243-E267, C273-P320, V339-V365, C376-S419, S455-K480, G501-T531, P605-G644, P657-M710, L721-M740, L741-A762, A763-P789, G797-W829, E830-V854, F958-Q1001, L34-G380 Tyrosine kinase catalytic domain signature: T713-R726, Y750-V768, I800-I810, S819-N841, C870-F892 Kinase receptor precursor: E31-C204 Ephrin receptor ligand binding domain: E31-C204 Signal peptide: M1-G30 Transmembrane region: V554-L561	MOTIFS HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
						SPScan HMMER HMMER

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4	7472006CD1	367	T310 T326 S349 S31 S158 S166 S290 S304		Protein kinases ATP-binding region signature: L18-K41 Serine/Threonine protein kinases active site signature: V132-L144 Eukaryotic protein kinase domain: Y12-M272 Testis specific serine/threonine kinase: M272-T364 Protein kinase domain: L14-I263 Tyrosine kinase catalytic domain signature: M90-K103, Y126-L144, Y197-S219	MOTIFS HMMR-PFAM BLAST- PRODOM BLAST-DOMO BLIMPS- PRINTS
5	2902460CD1	798	S56 S65 T67 T96 S98 T123 S132 S451 T428 S462 S463 Y464 S467 S473 T602 Y603 T634 T715 S18 S69 S116 S179 S292 S324 S386 S440 S499 S515 S531 S616	N317	Signal peptide: M1-A24 Calmodulin-binding domain: DM07435 P42335 210-672: P332-L797 Proline-rich protein: DM01369 B39066 172-256: G274-P330 1-D myoinositol tris-phosphate 3 kinase, EC 2.7.1.127, inositol 1,4,5-tris-phosphate, IP3K, IP3, transferase, kinase, calmodulin-binding: PD138098: G120-S510	SPScan MOTIFS BLAST- PRODOM BLAST-DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	6383934CD1	358	Y293 T48 S349 S31 S158 S258 S284 T340		Protein kinase ATP-binding domain: L18-K41 Protein kinase ST: I132-L144 Tyrosine kinase catalytic domain signature: M90-K103, Y126-L144, Y197-S219 Eukaryotic protein kinase domain: Y12-L272	MOTIFS PFAM BLIMPS-PRINTS
7	3210906CD1	1049	S306 S9 S111 T214 T346 S370 S375 T671 T701 S806 S853 S894 S1014 S60 S62 S453 T468 S521 T586 T604 T671 S742 T757 T776 T793 T886 S889 S910 T990 Y309	N1042	Protein kinase domain: DM00004 I48609 55-294: L18-R260 Testis specific serine/threonine kinase 2 protein kinase: PD029090: L272-T358 Protein kinase domain: DM00004 JC1446 20-261: V14-I263 Protein kinase domain: DM00004 P46549 32-279: D30-R269 Protein kinase ST: M147-L158 Eukaryotic protein kinase domain: F28-V281 Protein kinases signatures and profile: E127-N180 Serine/threonine protein kinase TAO1: E618-P777	BLAST- PRODOM BLAST-DOMO BLAST-DOMO MOTIFS HMMER-PFAM PROFILES SCAN BLAST- PRODOM MOTIFS
8	3339024CD1	322	S42 S117 T246 S266 S284 T109 T172 T195 S231 S236	N17 N87 N94 N112		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
9	4436929CD1	1212	S77 T187 S259 S608 S873 S9 S17 T59 S112 T124 T222 S264 T319 S324 S326 S548 S567 S604 S627 S680 S739 S740 T746 T747 S764 S778 T989 S1016 S1036 T1050 S1076 S255 S259 T309 T351 T557 T597 S604 S679 S687 S784 T869 S956 S1089 S1190 Y321 Y323 Y467	N33 N546 N624 N776 N1144	Eukaryotic protein kinase domain: F25-I289 Protein kinase domain: DM00004 P10676 18-272: L27-P278 CNH domain: Y894-R1192 Protein kinases signatures and profile: W129-T181 Protein kinase ST: V149-L161 NIK (Nck Interacting Kinase): PD147187: D496-W908	HMME-PFAM BLAST-DOMO HMME-PFAM PROFILES-SCAN MOTIFS BLAST-PRODROM
10	5046791CD1	280	S102 T161 Y162 T92 S209 S243 S102 T161	N155	Protein F55A11.6 C52E4.7, similar to Ser/Thr kinase: PD024191: G11-L130 Protein chromosome C34C6.5 C4A8.07C I sphingosine XII cosmid ORF: PD014044: H8-P97	BLAST-PRODROM BLAST-PRODROM
11	1416174CD1	114			Protein kinase ATP-binding domain: I32-M55 Protein kinase ST: I145-L157	MOTIFS MOTIFS
12	3244919CD1	375	S92 S276 T9 T48 T125 S295 T360 Y52	N338	Eukaryotic protein kinase domain: F26-Q278 Tyrosine kinase catalytic domain: PR00109: V103-Q116, Y139-L157 Protein kinase domain: DM00004 P54644 122-362: I28-S275 DM08046 P05986 1-397: S3-P305	HMME-PFAM BLIMPS-PRINTS BLAST-DOMO

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
13	058860CB1	1859	1-837, 1111-1198	60122573D4	1	491
				058860R6 (MUSCNOT01)	370	1005
				3011528F6 (MUSCNOT07)	852	1341
				3016678T6 (MUSCNOT07)	1299	1859
				3500745F6 (PROSTUT13)	1	456
14	2041716CB1	3501	1-2773	g4454511.v113.gs_3.nt.edit	22	884
				6063491H1 (BRAENOT02)	715	1093
				2190612F6 (THYRTUT03)	1072	1658
				70168906V1	1392	1989
				70164503V1	1840	2664
				70168645V1	2056	2696
				70167500V1	2541	3123
				1383374T6 (BRAITUT08)	2688	3255
				543319R6 (OVARNOT02)	2943	3501
				g5679461.v113.gs_2.edit	1	3039
15	7472005CB1	3039	1-557, 2741-3039, 824-1827	g5686590.v113.gs_5	1	1104
				70166939V1	3381	3916
				6882904J1 (BRAHTDR03)	1399	2005
				7117043H1 (BRAHNOE01)	614	1253
				7090661H1 (BRAUTDR03)	914	1492
				6811472J1 (SKIRNOR01)	2436	3034
				6882520J1 (BRAHTDR03)	1	639
				3753286H1 (BRAHDIT04)	3643	3939
				7029494H1 (BRAXTDR12)	1692	2288
				6911565J1 (PITUDIR01)	2169	2757
				7176637H1 (BRSTTMC01)	2766	3358
				2695922F6 (UTRSNOT12)	3114	3571
				g3873504.v113.gs_3.nt	73	1149
				2011686H1 (TESTNOT03)	665	858
				g2821547	972	1381
				6383934H1 (FIBRUNT02)	874	1176
				5281219H1 (TESTNON04)	1	239
16	7472006CB1	1104	823-1104			
17	2902460CB1	3939	1-1642, 2515-3100, 3766-3939			
18	6383934CB1	1381	1-359			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
19	3210906CB1	3904	3815-3904, 1-449, 901-1443, 1486-1805, 3039-3432	533823R6 (BRAINOT03) 1807122F6 (SINTNOT13) 4785178H1 (BRATNOT03) 1439938F6 (THYRNOT03) 2654018H1 (THYMNOT04) 713861X11 (PROSTUT01) 1416996X310D1 (BRAINOT12) 4326355F6 (TLYMUNT01) 2512189F6 (CONUTUT01) 273994R6 (PANCADIT03) 860975R6 (BRAITUT03) 273994T6 (PANCADIT03) 70774378V1 70772051V1 3339024F6 (SPLNNOT10) 70775014V1	3136 3432 2594 267 1336 1 2383 959 1593 666 2717 1947 446 787 1 1395	3683 3904 2860 769 1629 529 2783 1404 2079 1101 3224 2561 1101 1465 629 1987
20	3339024CB1	1987	1-125, 1987, 1955-1987, 1461-1493	2986160H1 (CARGDIT01) 1852144T6 (LUNGFET03) 3136101F6 (SMCCNOT01) SCLA03429V1 93327187_CD 2606210F6 (LUNGTUT07) 2827761F6 (TLYMNOT03) SZAU00120V1	1299 3362 555 2886 251 2654 633 1	1588 3925 1107 3425 3925 3136 1149 575
21	4436929CB1	3925	1431-2791, 1-956	3085382H1 (HEAONOT03) 2956512H1 (KIDNFET01) SCLA04243V1 1741505R6 (HIPONON01) 2805893F6 (BLADTUT08) 6390331H1 (BONRNOT01) 91512902 260140R6 (HNT2RAT01) 70495437V1	1427 3153 2312 1919 1085 1 637 586 213	1723 3430 2865 2435 1544 262 1210 1200 880
22	5046791CB1	1210	1-244			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
23	1416174CB1	1521	1-792, 876-975	3869131H1 (BMARNOT03)	1	231
				1416174H1 (BRAINOT12)	155	402
				2169725T6 (ENDCNOT03)	933	1504
				1683338F6 (PROSNOT15)	1030	1521
				1284949T6 (COLNNOT16)	871	1489
				3272203F6 (BRAINOT20)	269	898
24	3244919CB1	1640	919-1535	2287966H1 (BRAINON01)	1429	1640
				6307341H1 (NERDTN03)	440	1134
				7177378H1 (BRAXDIC01)	1	526
				70570341V1	1201	1588
				5372702H1 (BRAINOT22)	1428	1633
				70568614V1	677	1336

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
13	058860CB1	MUSCNOT07
14	2041716CB1	BRAXNOT03
17	2902460CB1	BRAGNON02
18	6383934CB1	FIBRUNT02
19	3210906CB1	BRAITUT03
20	3339024CB1	THYRNOT08
21	4436929CB1	ENDCNOT03
22	5046791CB1	BRABDIR01
23	1416174CB1	CARGDIT01
24	3244919CB1	BRAINOT21

Table 6

Library	Vector	Library Description
MUSCNOT07	pINCY	Library was constructed using 2 micrograms of polyA RNA isolated from muscle tissue removed from the forearm of a 38-year-old Caucasian female during a soft tissue excision. Pathology indicated the surgical margins of re-excision were free of tumor. Pathology for the matched tumor tissue indicated intramuscular hemangioma. Patient history included a normal delivery. Patient medications included melatonin, Valium, and Tylenol PM. Family history included breast cancer in the mother; and benign hypertension, cerebrovascular disease, colon cancer, and type II diabetes in the grandparent(s).
BRAXNOT03	pINCY	Library was constructed using 1.5 micrograms of polyA RNA isolated from sensory-motor cortex tissue removed from the brain of a 35-year-old Caucasian male who died from cardiac failure. Pathology indicated moderate leptomeningeal fibrosis and multiple microinfarctions of the cerebral neocortex. Grossly, the brain regions examined and cranial nerves were unremarkable, showing no evidence of atrophy. No atherosclerosis of the major vessels was noted. Microscopically, the cerebral hemisphere revealed moderate fibrosis of the leptomeninges with focal calcifications. There was evidence of shrunken and slightly eosinophilic pyramidal neurons throughout the cerebral hemispheres. There were also multiple small microscopic areas of cavitation with surrounding gliosis scattered throughout the cerebral cortex. Special stains with Bielschowsky silver, Klüver-Barrera, and Congo Red revealed no evidence of neurofibrillary tangles or diffuse anorectic amyloid plaques, demyelination, and cerebral amyloid angiopathy, respectively. Patient history included dilated cardiomyopathy, congestive heart failure, cardiomegaly, and an enlarged spleen and liver. Patient medications included simethicone, Lasix, Digoxin, Colace, Zantac, captopril, and Vasotec.

Table 6 cont.

Library	Vector	Library Description
BRAGNON02	pINCY	The library was constructed from a normalized substantia nigra tissue library constructed from 4.2x10 ⁷ independent clones. Starting RNA was made from RNA isolated from substantia nigra tissue removed from an 81-year-old Caucasian female who died from a hemorrhage and ruptured thoracic aorta due to atherosclerosis. Pathology indicated moderate atherosclerosis involving the internal carotids, bilaterally; microscopic infarcts of the frontal cortex and hippocampus; and scattered diffuse amyloid plaques and neurofibrillary tangles, consistent with age. Grossly, the leptomeninges showed only mild thickening and hyalinization along the superior sagittal sinus. The remainder of the leptomeninges was thin and contained some congested blood vessels. Mild atrophy was found mostly in the frontal poles and lobes, and temporal lobes, bilaterally. Microscopically, there were pairs of Alzheimer type II astrocytes within the deep layers of the neocortex. There was increased satellitosis around neurons in the deep gray matter in the middle frontal cortex. The amygdala contained rare diffuse plaques and neurofibrillary tangles. The posterior hippocampus contained a microscopic area of cystic cavitation with hemosiderin-laden macrophages surrounded by reactive gliosis. Patient history included sepsis, cholangitis, post-operative atelectasis, pneumonia CAD, cardiomegaly due to left ventricular hypertrophy, splenomegaly, arteriolonephrosclerosis, nodular colloid goiter, emphysema, CHF, hypothyroidism, and peripheral vascular disease. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
FIBRUNT02	pINCY	The library was constructed from polyA RNA isolated from an untreated MG-63 cell line derived from an osteosarcoma removed from a 14-year-old Caucasian male
BRABDIR01	pINCY	Library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident. Patient history included Huntington's disease, emphysema, and tobacco abuse.
BRAITUT03	PSPORT1	Library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 17-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a grade 4 fibrillary giant and small-cell astrocytoma. Family history included benign hypertension and cerebrovascular disease.
ENDCNOT03	pINCY	Library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.

Table 6 cont.

Library	Vector	Library Description
THYRNOT08	pINCY	Library was constructed using RNA isolated from the diseased left thyroid tissue removed from a 13-year-old Caucasian female during a complete thyroidectomy. Pathology indicated lymphocytic thyroiditis. Pathology for the matched tumor tissue indicated grade 1 papillary carcinoma. Multiple lymph nodes from the right, left, and midline section of the neck were negative for tumor. Fragments of the thymus were benign. Fibroadipose tissue was identified in the right inferior and superior parathyroid regions. Multiple lymph nodes (2 of 6) from the right side of the neck contained microscopic foci of metastatic papillary carcinoma. Patient history included attention deficit disorder with hyperactivity. Previous surgeries included an operative procedure on the external ear. Patient medications included Prozac. Family history included chronic obstructive asthma in the mother; alcohol abuse, benign hypertension, and depressive disorder in the grandparent(s); and attention deficit disorder with hyperactivity in the sibling(s).
BRAINOT21	pINCY	Library was constructed using RNA isolated from diseased brain tissue removed from the left frontal lobe of a 46-year-old Caucasian male during a lobectomy. Pathology indicated focal cortical and subcortical scarring of the left frontal lobe, characterized by cavitation and extensive reactive changes, including marked gliosis and hemosiderin deposition, consistent with a history of remote severe head trauma. GFAP was positive in astrocytes. The pattern of reactivity is that of reactive gliosis. Patient history included traumatic intracranial hemorrhage and brain injury with loss of consciousness following head trauma. Family history included cerebrovascular disease, cerebrovascular disease, and atherosclerotic coronary artery disease.
CARGDIT01	pINCY	Library was constructed using RNA isolated from diseased cartilage tissue. Patient history included osteoarthritis.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) <i>J. Mol. Biol.</i> 215:403-410; Altschul, S.F. et al. (1997) <i>Nucleic Acids Res.</i> 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less <i>Full Length sequences</i> : Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) <i>Proc. Natl. Acad. Sci. USA</i> 85:2444-2448; Pearson, W.R. (1990) <i>Methods Enzymol.</i> 183:63-98; and Smith, T.F. and M.S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489.	ESTs: fasta E value=1.06E-6 <i>Assembled ESTs</i> : fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less <i>Full Length sequences</i> : fastx score=100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) <i>Nucleic Acids Res.</i> 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) <i>Methods Enzymol.</i> 266:88-105; and Attwood, T.K. et al. (1997) <i>J. Chem. Inf. Comput. Sci.</i> 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) <i>J. Mol. Biol.</i> 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) <i>Nucleic Acids Res.</i> 26:320-322; Durbin, R. et al. (1998) <i>Our World View</i> , in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less <i>Signal peptide hits</i> : Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score = 1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score = 120 or greater; Match length = 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPSscan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score = 3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
 - 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-12,
 - b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-12,
 - c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12, and
 - 10 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-12.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-12.
- 15 3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
- 20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:13-24.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
- 25 7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
- 30 9. A method for producing a polypeptide of claim 1, the method comprising:
 - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 - 35 b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- 5 a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24,
 b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:13-24,
 c) a polynucleotide sequence complementary to a),
 d) a polynucleotide sequence complementary to b), and
10 e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

15 13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization
20 complex is formed between said probe and said target polynucleotide or fragments thereof, and
 b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

25 14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
30 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

35

17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-12.

18. A method for treating a disease or condition associated with decreased expression of functional PKIN, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional PKIN, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional PKIN, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a

compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 5 a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in
- 10 the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method

15 comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of
- 20 the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at
- 25 least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and
- 30 d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

<110> INCYTE GENOMICS, INC.
 YANG, Junming
 BAUGHN, Mariah R.
 BURFORD, Neil
 AU-YOUNG, Janice
 LU, Dyung Aina M.
 REDDY, Roopa
 YUE, Henry
 YAO, Monique G.
 LAL, Preeti
 KAHN, Farrah A.

<120> HUMAN KINASES

<130> PI-0002 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/172,066; 60/176,107; 60/176,107; 60/177,731

<151> 1999-12-23; 2000-01-14; 2000-01-14; 2000-01-21

<160> 24

<170> PERL Program

<210> 1

<211> 466

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 058860CD1

<400> 1

Met	Glu	Asp	Gly	Thr	Pro	Asn	Glu	His	Phe	Tyr	Thr	Pro	Thr	Glu
1				5					10					15
Glu	Arg	Gly	Ser	Ala	Tyr	Glu	Ile	Trp	Arg	Ser	Asp	Ser	Phe	Gly
				20					25					30
Thr	Pro	Asn	Glu	Ala	Ile	Glu	Pro	Lys	Asp	Asn	Glu	Met	Pro	Pro
				35					40					45
Ser	Phe	Ile	Glu	Pro	Leu	Thr	Lys	Arg	Lys	Val	Tyr	Glu	Asn	Thr
				50					55					60
Thr	Leu	Gly	Phe	Ile	Val	Glu	Val	Glu	Gly	Leu	Pro	Val	Pro	Gly
				65					70					75
Val	Lys	Trp	Tyr	Arg	Asn	Lys	Ser	Leu	Leu	Glu	Pro	Asp	Glu	Arg
				80					85					90
Ile	Lys	Met	Glu	Arg	Val	Gly	Asn	Val	Cys	Ser	Leu	Glu	Ile	Ser
				95					100					105
Asn	Ile	Gln	Lys	Gly	Glu	Gly	Gly	Glu	Tyr	Met	Cys	His	Ala	Val
				110					115					120
Asn	Ile	Ile	Gly	Glu	Ala	Lys	Ser	Phe	Ala	Asn	Val	Asp	Ile	Met
				125					130					135
Pro	Gln	Glu	Glu	Arg	Val	Val	Ala	Leu	Pro	Pro	Pro	Val	Thr	His
				140					145					150
Gln	His	Val	Met	Glu	Phe	Asp	Leu	Glu	His	Thr	Thr	Ser	Ser	Arg
				155					160					165
Thr	Pro	Ser	Pro	Gln	Glu	Ile	Val	Leu	Glu	Val	Glu	Leu	Ser	Glu
				170					175					180
Lys	Asp	Val	Lys	Glu	Phe	Glu	Lys	Gln	Val	Lys	Ile	Val	Thr	Val
				185					190					195
Pro	Glu	Phe	Thr	Pro	Asp	His	Lys	Ser	Met	Ile	Val	Ser	Leu	Asp
				200					205					210
Val	Leu	Pro	Phe	Asn	Phe	Val	Asp	Pro	Asn	Met	Asp	Ser	Arg	Glu
				215					220					225
Gly	Glu	Asp	Lys	Glu	Leu	Lys	Ile	Asp	Leu	Glu	Val	Phe	Glu	Met
				230					235					240

```

Pro Pro Arg Phe Ile Met Pro Ile Cys Asp Phe Lys Ile Pro Glu
245 250 255
Asn Ser Asp Ala Val Phe Lys Cys Ser Val Ile Gly Ile Pro Thr
260 265 270
Pro Glu Val Lys Trp Tyr Lys Glu Tyr Met Cys Ile Glu Pro Asp
275 280 285
Asn Ile Lys Tyr Val Ile Ser Glu Glu Lys Gly Ser His Thr Leu
290 295 300
Lys Ile Arg Asn Val Cys Leu Ser Asp Ser Ala Thr Tyr Arg Cys
305 310 315
Arg Ala Val Asn Cys Val Gly Glu Ala Ile Cys Arg Gly Phe Leu
320 325 330
Thr Met Gly Asp Ser Glu Ile Phe Ala Val Ile Ala Lys Lys Ser
335 340 345
Lys Val Thr Leu Ser Ser Leu Met Glu Glu Leu Val Leu Lys Ser
350 355 360
Asn Tyr Thr Asp Ser Phe Phe Glu Phe Gln Val Val Glu Gly Pro
365 370 375
Pro Arg Phe Ile Lys Gly Ile Ser Asp Cys Tyr Ala Pro Ile Gly
380 385 390
Thr Ala Ala Tyr Phe Gln Cys Leu Val Arg Gly Ser Pro Arg Pro
395 400 405
Thr Val Tyr Trp Tyr Lys Asp Gly Lys Leu Val Gln Gly Arg Arg
410 415 420
Phe Thr Val Glu Glu Ser Gly Thr Gly Phe His Asn Leu Phe Ile
425 430 435
Thr Ser Leu Val Lys Ser Asp Glu Gly Glu Tyr Arg Cys Val Ala
440 445 450
Thr Asn Lys Ser Gly Met Ala Glu Ser Phe Ala Ala Leu Thr Leu
455 460 465
Thr

```

<210> 2
 <211> 513
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2041716CD1

```

<400> 2
Met Glu Gly Gly Pro Ala Val Cys Cys Gln Asp Pro Arg Ala Glu
1 5 10 15
Leu Val Glu Arg Val Ala Ala Ile Asp Val Thr His Leu Glu Glu
20 25 30
Ala Asp Gly Gly Pro Glu Pro Thr Arg Asn Gly Val Asp Pro Pro
35 40 45
Pro Arg Ala Arg Ala Ala Ser Val Ile Pro Gly Ser Thr Ser Arg
50 55 60
Leu Leu Pro Ala Arg Pro Ser Leu Ser Ala Arg Lys Leu Ser Leu
65 70 75
Gln Glu Arg Pro Ala Gly Ser Tyr Leu Glu Ala Gln Ala Gly Pro
80 85 90
Tyr Ala Thr Gly Pro Ala Ser His Ile Ser Pro Arg Ala Trp Arg
95 100 105
Arg Pro Thr Ile Glu Ser His His Val Ala Ile Ser Asp Ala Glu
110 115 120
Asp Cys Val Gln Leu Asn Gln Tyr Lys Leu Gln Ser Glu Ile Gly
125 130 135
Lys Val Gly Leu Thr Asp Ala Tyr Leu Gln Gly Ala Tyr Gly Val
140 145 150
Val Arg Leu Ala Tyr Asn Glu Ser Glu Asp Arg His Tyr Ala Met
155 160 165
Lys Val Leu Ser Lys Lys Lys Leu Leu Lys Gln Tyr Gly Phe Pro
170 175 180

```



```

Arg Arg Pro Pro Pro Arg Gly Ser Gln Ala Ala Gln Gly Gly Pro
185 190 195
Ala Lys Gln Leu Leu Pro Leu Glu Arg Val Tyr Gln Glu Ile Ala
200 205 210
Ile Leu Lys Lys Leu Asp His Val Asn Val Val Lys Leu Ile Glu
215 220 225
Val Leu Asp Asp Pro Ala Glu Asp Asn Leu Tyr Leu Val Asp Leu
230 235 240
Leu Arg Lys Gly Pro Val Met Glu Val Pro Cys Asp Lys Pro Phe
245 250 255
Ser Glu Glu Gln Ala Arg Leu Tyr Leu Arg Asp Val Ile Leu Gly
260 265 270
Leu Glu Tyr Leu His Cys Gln Lys Ile Val His Arg Asp Ile Lys
275 280 285
Pro Ser Asn Leu Leu Leu Gly Asp Asp Gly His Val Lys Ile Ala
290 295 300
Asp Phe Gly Val Ser Asn Gln Phe Glu Gly Asn Asp Ala Gln Leu
305 310 315
Ser Ser Thr Ala Gly Thr Pro Ala Phe Met Ala Pro Glu Ala Ile
320 325 330
Ser Asp Ser Gly Gln Ser Phe Ser Gly Lys Ala Leu Asp Val Trp
335 340 345
Ala Thr Gly Val Thr Leu Tyr Cys Phe Val Tyr Gly Lys Cys Pro
350 355 360
Phe Ile Asp Asp Phe Ile Leu Ala Leu His Arg Lys Ile Lys Asn
365 370 375
Glu Pro Val Val Phe Pro Glu Glu Pro Glu Ile Ser Glu Glu Leu
380 385 390
Lys Asp Leu Ile Leu Lys Met Leu Asp Lys Asn Pro Glu Thr Arg
395 400 405
Ile Gly Val Pro Asp Ile Lys Leu His Pro Trp Val Thr Lys Asn
410 415 420
Gly Glu Glu Pro Leu Pro Ser Glu Glu Glu His Cys Ser Val Val
425 430 435
Glu Val Thr Glu Glu Glu Val Lys Asn Ser Val Arg Leu Ile Pro
440 445 450
Ser Trp Thr Thr Val Ile Leu Val Lys Ser Met Leu Arg Lys Arg
455 460 465
Ser Phe Gly Asn Pro Phe Glu Pro Gln Ala Arg Arg Glu Glu Arg
470 475 480
Ser Met Ser Ala Pro Gly Asn Leu Leu Val Lys Glu Gly Phe Gly
485 490 495
Glu Gly Gly Lys Ser Pro Glu Leu Pro Gly Val Gln Glu Asp Glu
500 505 510
Ala Ala Ser

```

<210> 3
 <211> 1012
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7472005CD1

```

<400> 3
Met Ala Pro Ala Arg Gly Arg Leu Pro Pro Ala Leu Trp Val Val
1 5 10 15
Thr Ala Ala Ala Ala Ala Thr Cys Val Ser Ala Ala Arg Gly
20 25 30
Glu Val Asn Leu Leu Asp Thr Ser Thr Ile His Gly Asp Trp Gly
35 40 45
Trp Leu Thr Tyr Pro Ala His Gly Trp Asp Ser Ile Asn Glu Val
50 55 60
Asp Glu Ser Phe Gln Pro Ile His Thr Tyr Gln Val Cys Asn Val
65 70 75

```

Met	Ser	Pro	Asn	Gln	Asn	Asn	Trp	Leu	Arg	Thr	Ser	Trp	Val	Pro
				80					85					90
Arg	Asp	Gly	Ala	Arg	Arg	Val	Tyr	Ala	Glu	Ile	Lys	Phe	Thr	Leu
				95					100					105
Arg	Asp	Cys	Asn	Ser	Met	Pro	Gly	Val	Leu	Gly	Thr	Cys	Lys	Glu
				110					115					120
Thr	Phe	Asn	Leu	Tyr	Tyr	Leu	Glu	Ser	Asp	Arg	Asp	Leu	Gly	Ala
				125					130					135
Ser	Thr	Gln	Glu	Ser	Gln	Phe	Leu	Lys	Ile	Asp	Thr	Ile	Ala	Ala
				140					145					150
Asp	Glu	Ser	Phe	Thr	Gly	Ala	Asp	Leu	Gly	Val	Arg	Arg	Leu	Lys
				155					160					165
Leu	Asn	Thr	Glu	Val	Arg	Ser	Val	Gly	Pro	Leu	Ser	Lys	Arg	Gly
				170					175					180
Phe	Tyr	Leu	Ala	Phe	Gln	Asp	Ile	Gly	Ala	Cys	Leu	Ala	Ile	Leu
				185					190					195
Ser	Leu	Arg	Ile	Tyr	Lys	Lys	Cys	Pro	Ala	Met	Val	Arg	Asn	
				200					205					210
Leu	Ala	Ala	Phe	Ser	Glu	Ala	Val	Thr	Gly	Ala	Asp	Ser	Ser	Ser
				215					220					225
Leu	Val	Glu	Val	Arg	Gly	Gln	Cys	Val	Arg	His	Ser	Glu	Glu	Arg
				230					235					240
Asp	Thr	Pro	Lys	Met	Tyr	Cys	Ser	Ala	Glu	Gly	Glu	Trp	Leu	Val
				245					250					255
Pro	Ile	Gly	Lys	Cys	Val	Cys	Ser	Ala	Gly	Tyr	Glu	Glu	Arg	Arg
				260					265					270
Asp	Ala	Cys	Val	Ala	Cys	Glu	Leu	Gly	Phe	Tyr	Lys	Ser	Ala	Pro
				275					280					285
Gly	Asp	Gln	Leu	Cys	Ala	Arg	Cys	Pro	Pro	His	Ser	His	Ser	Ala
				290					295					300
Ala	Pro	Ala	Ala	Gln	Ala	Cys	His	Cys	Asp	Leu	Ser	Tyr	Tyr	Arg
				305					310					315
Ala	Ala	Leu	Asp	Pro	Pro	Ser	Ser	Ala	Cys	Thr	Arg	Pro	Pro	Ser
				320					325					330
Ala	Pro	Val	Asn	Leu	Ile	Ser	Ser	Val	Asn	Gly	Thr	Ser	Val	Thr
				335					340					345
Leu	Glu	Trp	Ala	Pro	Pro	Leu	Asp	Pro	Gly	Gly	Arg	Ser	Asp	Ile
				350					355					360
Thr	Tyr	Asn	Ala	Val	Cys	Arg	Arg	Cys	Pro	Trp	Ala	Leu	Ser	Arg
				365					370					375
Cys	Glu	Ala	Cys	Gly	Ser	Gly	Thr	Arg	Phe	Val	Pro	Gln	Gln	Thr
				380					385					390
Ser	Leu	Val	Gln	Ala	Ser	Leu	Leu	Val	Ala	Asn	Leu	Leu	Ala	His
				395					400					405
Met	Asn	Tyr	Ser	Phe	Trp	Ile	Glu	Ala	Val	Asn	Gly	Val	Ser	Asp
				410					415					420
Leu	Ser	Pro	Glu	Pro	Arg	Arg	Ala	Ala	Val	Val	Asn	Ile	Thr	Thr
				425					430					435
Asn	Gln	Ala	Ala	Pro	Ser	Gln	Val	Val	Val	Ile	Arg	Gln	Glu	Arg
				440					445					450
Ala	Gly	Gln	Thr	Ser	Val	Ser	Leu	Leu	Trp	Gln	Glu	Pro	Glu	Gln
				455					460					465
Pro	Asn	Gly	Ile	Ile	Leu	Glu	Tyr	Glu	Ile	Lys	Tyr	Tyr	Glu	Lys
				470					475					480
Asp	Lys	Glu	Met	Gln	Ser	Tyr	Ser	Thr	Leu	Lys	Ala	Val	Thr	Thr
				485					490					495
Arg	Ala	Thr	Val	Ser	Gly	Leu	Lys	Pro	Gly	Thr	Arg	Tyr	Val	Phe
				500					505					510
Gln	Val	Arg	Ala	Arg	Thr	Ser	Ala	Gly	Cys	Gly	Arg	Phe	Ser	Gln
				515					520					525
Ala	Met	Glu	Val	Glu	Thr	Gly	Lys	Pro	Arg	Pro	Arg	Tyr	Asp	Thr
				530					535					540
Arg	Thr	Ile	Val	Trp	Ile	Cys	Leu	Thr	Leu	Ile	Thr	Gly	Leu	Val
				545					550					555
Val	Leu	Leu	Leu	Leu	Leu	Ile	Cys	Lys	Lys	Arg	His	Cys	Gly	Tyr
				560					565					570
Ser	Lys	Ala	Phe	Gln	Asp	Ser	Asp	Glu	Glu	Lys	Met	His	Tyr	Gln

Asn Gly Gln Ala	575	580	585
Pro Pro Pro Val Phe	590	595	600
Pro Gly Lys Leu	605	610	615
Tyr Glu Glu Pro	620	625	630
Glu Ala Ser Arg	635	640	645
Ser Gly Glu Val	650	655	660
Asp Val Pro Val	665	670	675
Arg Gln Arg Arg	680	685	690
Phe Asp His Pro	695	700	705
Gly Arg Leu Ala	710	715	720
Leu Asp Thr Phe	725	730	735
Gln Leu Val Gly	740	745	750
Leu Ser Asp Leu	755	760	765
Val Leu Val Asp	770	775	780
Leu Ser Arg Val	785	790	795
Thr Gly Gly Lys	800	805	810
Ala Phe Arg Thr	815	820	825
Val Val Met Trp	830	835	840
Asn Met Thr Asn	845	850	855
Ser Ser Val Glu	860	865	870
Pro His Ala Leu	875	880	885
Arg Ala Gln Arg	890	895	900
Ala Leu Ile Arg	905	910	915
Ser Arg Cys Pro	920	925	930
Arg Gly Gly Ser	935	940	945
Leu Asp Ser Ile	950	955	960
Gly Gly Tyr Ser	965	970	975
Asp Val Arg Ala	980	985	990
Ile Leu Gly Ser	995	1000	1005
Gln Gly Pro Arg	1010		

<210> 4

<211> 367

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472006CD1

<400> 4

```

Met Asp Asp Ala Ala Val Leu Lys Arg Arg Gly Tyr Leu Leu Gly
 1          5          10          15
Ile Asn Leu Gly Glu Gly Ser Tyr Ala Lys Val Lys Ser Ala Tyr
          20          25          30
Ser Glu Arg Leu Lys Phe Asn Val Ala Ile Lys Ile Ile Asp Arg
          35          40          45
Lys Lys Ala Pro Ala Asp Phe Leu Glu Lys Phe Leu Pro Arg Glu
          50          55          60
Ile Glu Ile Leu Ala Met Leu Asn His Cys Ser Ile Ile Lys Thr
          65          70          75
Tyr Glu Ile Phe Glu Thr Ser His Gly Lys Val Tyr Ile Val Met
          80          85          90
Glu Leu Ala Val Gln Gly Asp Leu Leu Glu Leu Ile Lys Thr Arg
          95          100          105
Gly Ala Leu His Glu Asp Glu Ala Arg Lys Lys Phe His Gln Leu
          110          115          120
Ser Leu Ala Ile Lys Tyr Cys His Asp Leu Asp Val Val His Arg
          125          130          135
Asp Leu Lys Cys Asp Asn Leu Leu Leu Asp Lys Asp Phe Asn Ile
          140          145          150
Lys Leu Ser Asp Phe Ser Phe Ser Lys Arg Cys Leu Arg Asp Asp
          155          160          165
Ser Gly Arg Met Ala Leu Ser Lys Thr Phe Cys Gly Ser Pro Ala
          170          175          180
Tyr Ala Ala Pro Glu Val Leu Gln Gly Ile Pro Tyr Gln Pro Lys
          185          190          195
Val Tyr Asp Ile Trp Ser Leu Gly Val Ile Leu Tyr Ile Met Val
          200          205          210
Cys Gly Ser Met Pro Tyr Asp Asp Ser Asn Ile Lys Lys Met Leu
          215          220          225
Arg Ile Gln Lys Glu His Arg Val Asn Phe Pro Arg Ser Lys His
          230          235          240
Leu Thr Gly Glu Cys Lys Asp Leu Ile Tyr His Met Leu Gln Pro
          245          250          255
Asp Val Asn Arg Arg Leu His Ile Asp Glu Ile Leu Ser His Cys
          260          265          270
Trp Met Gln Pro Lys Ala Arg Gly Ser Pro Ser Val Ala Ile Asn
          275          280          285
Lys Glu Gly Glu Ser Ser Arg Gly Thr Glu Pro Leu Trp Thr Pro
          290          295          300
Glu Pro Gly Ser Asp Lys Lys Ser Ala Thr Lys Leu Glu Pro Glu
          305          310          315
Gly Glu Ala Gln Pro Gln Ala Gln Pro Glu Thr Lys Pro Glu Gly
          320          325          330
Thr Ala Met Gln Met Ser Arg Gln Ser Glu Ile Leu Gly Phe Pro
          335          340          345
Ser Lys Pro Ser Thr Met Glu Thr Glu Glu Gly Pro Pro Gln Gln
          350          355          360
Pro Pro Glu Thr Arg Ala Gln
          365

```

<210> 5

<211> 798

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2902460CD1

<400> 5

```

Met Phe Glu Ala His Ile Gln Ala Gln Ser Ser Ala Ile Gln Ala
 1          5          10          15
Pro Arg Ser Pro Arg Leu Gly Arg Ala Arg Ser Pro Ser Pro Cys

```

Pro Phe Arg Ser	20	25	30
Ser Ser Gln Pro Pro Gly Arg Val Leu Val Gln	35	40	45
Gly Ala Arg Ser Glu Glu Arg Arg Thr Lys Ser Trp Gly Glu Gln	50	55	60
Cys Pro Glu Thr Ser Gly Thr Asp Ser Gly Arg Lys Gly Gly Pro	65	70	75
Ser Leu Cys Ser Ser Gln Val Lys Lys Gly Met Pro Pro Leu Pro	80	85	90
Gly Arg Ala Ala Pro Thr Gly Ser Glu Ala Gln Gly Pro Ser Ala	95	100	105
Phe Val Arg Met Glu Lys Gly Ile Pro Ala Ser Pro Arg Cys Gly	110	115	120
Ser Pro Thr Ala Met Glu Ile Asp Lys Arg Gly Ser Pro Thr Pro	125	130	135
Gly Thr Arg Ser Cys Leu Ala Pro Ser Leu Gly Leu Phe Gly Ala	140	145	150
Ser Leu Thr Met Ala Thr Glu Val Ala Ala Arg Val Thr Ser Thr	155	160	165
Gly Pro His Arg Pro Gln Asp Leu Ala Leu Thr Glu Pro Ser Gly	170	175	180
Arg Ala Arg Glu Leu Glu Asp Leu Gln Pro Pro Glu Ala Leu Val	185	190	195
Glu Arg Gln Gly Gln Phe Leu Gly Ser Glu Thr Ser Pro Ala Pro	200	205	210
Glu Arg Gly Gly Pro Arg Asp Gly Glu Pro Pro Gly Lys Met Gly	215	220	225
Lys Gly Tyr Leu Pro Cys Gly Met Pro Gly Ser Gly Glu Pro Glu	230	235	240
Val Gly Lys Arg Pro Glu Glu Thr Thr Val Ser Val Gln Ser Ala	245	250	255
Glu Ser Ser Asp Ala Leu Ser Trp Ser Arg Leu Pro Arg Ala Leu	260	265	270
Ala Ser Val Gly Pro Glu Glu Ala Arg Ser Gly Ala Pro Val Gly	275	280	285
Gly Gly Arg Trp Gln Leu Ser Asp Arg Val Glu Gly Gly Ser Pro	290	295	300
Thr Leu Gly Leu Leu Gly Gly Ser Pro Ser Ala Gln Pro Gly Thr	305	310	315
Gly Asn Val Glu Ala Gly Ile Pro Ser Gly Arg Met Leu Glu Pro	320	325	330
Leu Pro Cys Trp Asp Ala Ala Lys Asp Leu Lys Glu Pro Gln Cys	335	340	345
Pro Pro Gly Asp Arg Val Gly Val Gln Pro Gly Asn Ser Arg Val	350	355	360
Trp Gln Gly Thr Met Glu Lys Ala Gly Leu Ala Trp Thr Arg Gly	365	370	375
Thr Gly Val Gln Ser Glu Gly Thr Trp Glu Ser Gln Arg Gln Asp	380	385	390
Ser Asp Ala Leu Pro Ser Pro Glu Leu Leu Pro Gln Asp Gln Asp	395	400	405
Lys Pro Phe Leu Arg Lys Ala Cys Ser Pro Ser Asn Ile Pro Ala	410	415	420
Val Ile Ile Thr Asp Met Gly Thr Gln Glu Asp Gly Ala Leu Glu	425	430	435
Glu Thr Gln Gly Ser Pro Arg Gly Asn Leu Pro Leu Arg Lys Leu	440	445	450
Ser Ser Ser Ser Ala Ser Ser Thr Gly Phe Ser Ser Ser Tyr Glu	455	460	465
Asp Ser Glu Glu Asp Ile Ser Ser Asp Pro Glu Arg Thr Leu Asp	470	475	480
Pro Asn Ser Ala Phe Leu His Thr Leu Asp Gln Gln Lys Pro Arg	485	490	495
Val Ser Lys Ser Trp Arg Lys Ile Lys Asn Met Val His Trp Ser	500	505	510
Pro Phe Val Met Ser Phe Lys Lys Lys Tyr Pro Trp Ile Gln Leu	515	520	525

Ala Gly His Ala Gly Ser Phe Lys Ala Ala Ala Asn Gly Arg Ile	530	535	540
Leu Lys Lys His Cys Glu Ser Glu Gln Arg Cys Leu Asp Arg Leu	545	550	555
Met Val Asp Val Leu Arg Pro Phe Val Pro Ala Tyr His Gly Asp	560	565	570
Val Val Lys Asp Gly Glu Arg Tyr Asn Gln Met Asp Asp Leu Leu	575	580	585
Ala Asp Phe Asp Ser Pro Cys Val Met Asp Cys Lys Met Gly Ile	590	595	600
Arg Thr Tyr Leu Glu Glu Glu Leu Thr Lys Ala Arg Lys Lys Pro	605	610	615
Ser Leu Arg Lys Asp Met Tyr Gln Lys Met Ile Glu Val Asp Pro	620	625	630
Glu Ala Pro Thr Glu Glu Glu Lys Ala Gln Arg Ala Val Thr Lys	635	640	645
Pro Arg Tyr Met Gln Trp Arg Glu Thr Ile Ser Ser Thr Ala Thr	650	655	660
Leu Gly Phe Arg Ile Glu Gly Ile Lys Lys Glu Asp Gly Thr Val	665	670	675
Asn Arg Asp Phe Lys Lys Thr Lys Thr Arg Glu Gln Val Thr Glu	680	685	690
Ala Phe Arg Glu Phe Thr Lys Gly Asn His Asn Ile Leu Ile Ala	695	700	705
Tyr Arg Asp Arg Leu Lys Ala Ile Arg Thr Thr Leu Glu Val Ser	710	715	720
Pro Phe Phe Lys Cys His Glu Val Ile Gly Ser Ser Leu Leu Phe	725	730	735
Ile His Asp Lys Lys Glu Gln Ala Lys Val Trp Met Ile Asp Phe	740	745	750
Gly Lys Thr Thr Pro Leu Pro Glu Gly Gln Thr Leu Gln His Asp	755	760	765
Val Pro Trp Gln Glu Gly Asn Arg Glu Asp Gly Tyr Leu Ser Gly	770	775	780
Leu Asn Asn Leu Val Asp Ile Leu Thr Glu Met Ser Gln Asp Ala	785	790	795
Pro Leu Ala			

<210> 6
 <211> 358
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6383934CD1

<400> 6	
Met Asp Asp Ala Thr Val Leu Arg Lys Lys Gly Tyr Ile Val Gly	15
1 5 10	
Ile Asn Leu Gly Lys Gly Ser Tyr Ala Lys Val Lys Ser Ala Tyr	30
20 25	
Ser Glu Arg Leu Lys Phe Asn Val Ala Val Lys Ile Ile Asp Arg	45
35 40	
Lys Lys Thr Pro Thr Asp Phe Val Glu Arg Phe Leu Pro Arg Glu	60
50 55	
Met Asp Ile Leu Ala Thr Val Asn His Gly Ser Ile Ile Lys Thr	75
65 70	
Tyr Glu Ile Phe Glu Thr Ser Asp Gly Arg Ile Tyr Ile Ile Met	90
80 85	
Glu Leu Gly Val Gln Gly Asp Leu Leu Glu Phe Ile Lys Cys Gln	105
95 100	
Gly Ala Leu His Glu Asp Val Ala Arg Lys Met Phe Arg Gln Leu	120
110 115	
Ser Ser Ala Val Lys Tyr Cys His Asp Leu Asp Ile Val His Arg	135
125 130	

```

Asp Leu Lys Cys Glu Asn Leu Leu Leu Asp Lys Asp Phe Asn Ile
140 145 150
Lys Leu Ser Asp Phe Gly Phe Ser Lys Arg Cys Leu Arg Asp Ser
155 160 165
Asn Gly Arg Ile Ile Leu Ser Lys Thr Phe Cys Gly Ser Ala Ala
170 175 180
Tyr Ala Ala Pro Glu Val Leu Gln Ser Ile Pro Tyr Gln Pro Lys
185 190 195
Val Tyr Asp Ile Trp Ser Leu Gly Val Ile Leu Tyr Ile Met Val
200 205 210
Cys Gly Ser Met Pro Tyr Asp Asp Ser Asp Ile Lys Lys Met Leu
215 220 225
Arg Ile Gln Lys Glu His Arg Val Asn Phe Pro Arg Ser Lys His
230 235 240
Leu Thr Cys Glu Cys Lys Asp Leu Ile Tyr His Met Leu Gln Pro
245 250 255
Asp Val Ser Gln Arg Leu His Ile Asp Glu Ile Leu Ser His Ser
260 265 270
Trp Leu Gln Pro Pro Lys Pro Lys Ala Thr Ser Ser Ala Ser Phe
275 280 285
Lys Arg Glu Gly Glu Gly Lys Tyr Arg Ala Glu Cys Lys Leu Asp
290 295 300
Thr Lys Thr Gly Leu Arg Pro Asp His Arg Pro Asp His Lys Leu
305 310 315
Gly Ala Lys Thr Gln His Arg Leu Leu Val Val Pro Glu Asn Glu
320 325 330
Asn Arg Met Glu Asp Arg Leu Ala Glu Thr Ser Arg Ala Lys Asp
335 340 345
His His Ile Ser Gly Ala Glu Val Gly Lys Ala Ser Thr
350 355

```

<210> 7

<211> 1049

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3210906CD1

<400> 7

```

Met Pro Ala Gly Gly Arg Ala Gly Ser Leu Lys Asp Pro Asp Val
1 5 10 15
Ala Glu Leu Phe Phe Lys Asp Asp Pro Glu Lys Leu Phe Ser Asp
20 25 30
Leu Arg Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe Ala
35 40 45
Arg Asp Val Arg Asn Ser Glu Val Val Ala Ile Lys Lys Met Ser
50 55 60
Tyr Ser Gly Lys Gln Ser Asn Glu Lys Trp Gln Asp Ile Ile Lys
65 70 75
Glu Val Arg Phe Leu Gln Lys Leu Arg His Pro Asn Thr Ile Gln
80 85 90
Tyr Arg Gly Cys Tyr Leu Arg Glu His Thr Ala Trp Leu Val Met
95 100 105
Glu Tyr Cys Leu Gly Ser Thr Ser Asp Leu Leu Glu Val His Lys
110 115 120
Lys Pro Leu Gln Glu Val Glu Ile Ala Ala Val Thr His Gly Ala
125 130 135
Leu Gln Gly Leu Ala Tyr Leu His Ser His Asn Met Ile His Arg
140 145 150
Asp Val Lys Ala Gly Asn Ile Leu Leu Ser Glu Pro Gly Leu Val
155 160 165
Lys Leu Gly Asp Phe Gly Ser Ala Ser Ile Met Ala Pro Ala Asn
170 175 180
Ser Phe Val Gly Thr Pro Tyr Trp Met Ala Pro Glu Val Ile Leu
185 190 195

```

Ala Met Asp Glu Gly Gln Tyr Asp Gly Lys Val Asp Val Trp Ser	200	205	210
Leu Gly Ile Thr Cys Ile Glu Leu Ala Glu Arg Lys Pro Pro Leu	215	220	225
Phe Asn Met Asn Ala Met Ser Ala Leu Tyr His Ile Ala Gln Asn	230	235	240
Glu Ser Pro Val Leu Gln Ser Gly His Trp Ser Glu Tyr Phe Arg	245	250	255
Asn Phe Val Asp Ser Cys Leu Gln Lys Ile Pro Gln Asp Arg Pro	260	265	270
Thr Ser Glu Val Leu Leu Lys His Arg Phe Val Leu Arg Glu Arg	275	280	285
Pro Pro Thr Val Ile Met Asp Leu Ile Gln Arg Thr Lys Asp Ala	290	295	300
Val Arg Glu Leu Asp Ser Leu Gln Tyr Arg Lys Met Lys Lys Ile	305	310	315
Leu Phe Gln Glu Ala Pro Asn Gly Pro Gly Ala Glu Ala Pro Glu	320	325	330
Glu Glu Glu Glu Ala Glu Pro Tyr Met His Leu Ala Gly Thr Leu	335	340	345
Thr Ser Leu Glu Ser Ser His Ser Val Pro Ser Met Ser Ile Ser	350	355	360
Ala Ser Ser Gln Ser Ser Ser Val Asn Ser Leu Ala Asp Ala Ser	365	370	375
Asp Asn Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu	380	385	390
Glu Glu Gly Pro Glu Ala Arg Glu Met Ala Met Met Gln Glu Gly	395	400	405
Glu His Thr Val Thr Ser His Ser Ser Ile Ile His Arg Leu Pro	410	415	420
Gly Ser Asp Asn Leu Tyr Asp Asp Pro Tyr Gln Pro Glu Ile Thr	425	430	435
Pro Ser Pro Leu Gln Pro Pro Ala Ala Pro Ala Pro Thr Ser Thr	440	445	450
Thr Ser Ser Ala Arg Arg Arg Ala Tyr Cys Arg Asn Arg Asp His	455	460	465
Phe Ala Thr Ile Arg Thr Ala Ser Leu Val Ser Arg Gln Ile Gln	470	475	480
Glu His Glu Gln Asp Ser Ala Leu Arg Glu Gln Leu Ser Gly Tyr	485	490	495
Lys Arg Met Arg Arg Gln His Gln Lys Gln Leu Leu Ala Leu Glu	500	505	510
Ser Arg Leu Arg Gly Glu Arg Glu Glu His Ser Ala Arg Leu Gln	515	520	525
Arg Glu Leu Glu Ala Gln Arg Ala Gly Phe Gly Ala Glu Ala Glu	530	535	540
Lys Leu Ala Arg Arg His Gln Ala Ile Gly Glu Lys Glu Ala Arg	545	550	555
Ala Ala Gln Ala Glu Glu Arg Lys Phe Gln Gln His Ile Leu Gly	560	565	570
Gln Gln Lys Lys Glu Leu Ala Ala Leu Leu Glu Ala Gln Lys Arg	575	580	585
Thr Tyr Lys Leu Arg Lys Glu Gln Leu Lys Glu Glu Leu Gln Glu	590	595	600
Asn Pro Ser Thr Pro Lys Arg Glu Lys Ala Glu Trp Leu Leu Arg	605	610	615
Gln Lys Glu Gln Leu Gln Gln Cys Gln Ala Glu Glu Glu Ala Gly	620	625	630
Leu Leu Arg Arg Gln Arg Gln Tyr Phe Glu Leu Gln Cys Arg Gln	635	640	645
Tyr Lys Arg Lys Met Leu Leu Ala Arg His Ser Leu Asp Gln Asp	650	655	660
Leu Leu Arg Glu Asp Leu Asn Lys Lys Gln Thr Gln Lys Asp Leu	665	670	675
Glu Cys Ala Leu Leu Leu Arg Gln His Glu Ala Thr Arg Glu Leu	680	685	690
Glu Leu Arg Gln Leu Gln Ala Val Gln Arg Thr Arg Ala Glu Leu			

Thr Arg Leu Gln	695	700	705
His Gln Thr Glu Leu	710	715	720
Asn Lys Arg Arg	725	730	735
Val Arg Gln Gln	740	745	750
Lys Lys Gln Phe	755	760	765
Lys Ala Leu Arg	770	775	780
His Lys Ser Leu	785	790	795
Leu Ala Ile Leu	800	805	810
Leu Ser Ser Gln	815	820	825
Phe Gln Ala Leu	830	835	840
Asn Ala Tyr Gln	845	850	855
Glu Arg Glu Leu	860	865	870
Ala Leu Leu Glu	875	880	885
Thr Gly Arg Ser	890	895	900
Arg Glu Ile Glu	905	910	915
Ser Ser Met Ala	920	925	930
Gly Tyr Pro Ala	935	940	945
Val Pro Arg Ser	950	955	960
Gly Met Pro Pro	965	970	975
Pro Gly Pro Pro	980	985	990
Pro Arg Gly Gly	995	1000	1005
Leu Arg Arg Ala	1010	1015	1020
Pro Pro Ala Ala	1025	1030	1035
Val Ala Ser His	1040	1045	

<210> 8

<211> 322

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3339024CD1

<400> 8

Met Pro Thr Phe Ser	Ile Pro Gly Thr	Leu Glu Ser Gly His	Pro
1	5	10	15
Arg Asn Leu Thr Cys	Ser Val Pro Trp	Ala Cys Glu Gln Gly	Thr
20	25	30	
Pro Pro Thr Ile Thr	Trp Met Gly Ala	Ser Val Ser Ser Leu	Asp
35	40	45	
Pro Thr Ile Thr Arg	Ser Ser Met Leu	Ser Leu Ile Pro Gln	Pro
50	55	60	
Gln Asp His Gly Thr	Ser Leu Thr Cys	Gln Val Thr Leu Pro	Gly

Ala Gly Val Thr	Met	Thr Arg Ala Val	Arg	Leu Asn Ile Ser	Tyr
Pro Pro Gln Asn	Leu	Thr Met Thr Val	Phe	Gln Gly Asp Gly	Thr
Ala Ser Thr Thr	Leu	Arg Asn Gly Ser	Ala	Leu Ser Val Leu	Glu
Gly Gln Ser Leu	His	Leu Val Cys Ala	Val	Asp Ser Asn Pro	Pro
Ala Arg Leu Ser	Trp	Thr Trp Gly Ser	Leu	Thr Leu Ser Pro	Ser
Gln Ser Ser Asn	Leu	Gly Val Leu Glu	Leu	Pro Arg Val His	Val
Lys Asp Glu Gly	Glu	Phe Thr Cys Arg	Ala	Gln Asn Pro Leu	Gly
Ser Gln His Ile	Ser	Leu Ser Leu Ser	Leu	Gln Asn Glu Tyr	Thr
Gly Lys Met Arg	Pro	Ile Ser Gly Val	Thr	Leu Gly Ala Phe	Gly
Gly Ala Gly Ala	Thr	Ala Leu Val Phe	Leu	Tyr Phe Cys Ile	Ile
Phe Val Val Val	Arg	Ser Cys Arg Lys	Lys	Ser Ala Arg Pro	Ala
Val Gly Val Gly	Asp	Thr Gly Met Glu	Asp	Ala Asn Ala Val	Trp
Gly Ser Ala Ser	Gln	Gly Pro Leu Ile	Glu	Ser Pro Ala Asp	Asp
Ser Pro Pro His	His	Ala Pro Pro Ala	Leu	Ala Thr Pro Ser	Pro
Glu Glu Gly Glu	Ile	Gln Tyr Ala Ser	Leu	Ser Phe His Lys	Ala
Arg Pro Gln Tyr	Pro	Gln Glu Gln Glu	Ala	Ile Gly Tyr Glu	Tyr
Ser Glu Ile Asn	Ile	Pro Lys			

<210> 9
 <211> 1212
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4436929CD1

Met Ala Asn Asp	Ser	Pro Ala Lys Ser	Leu	Val Asp Ile Asp	Leu
1	5		10		15
Ser Ser Leu Arg	Asp	Pro Ala Gly Ile	Phe	Glu Leu Val Glu	Val
	20		25		30
Val Gly Asn Gly	Thr	Tyr Gly Gln Val	Tyr	Lys Gly Arg His	Val
	35		40		45
Lys Thr Gly Gln	Leu	Ala Ala Ile Lys	Val	Met Asp Val Thr	Glu
	50		55		60
Asp Glu Glu Glu	Glu	Ile Lys Leu Glu	Ile	Asn Met Leu Lys	Lys
	65		70		75
Tyr Ser His His	Arg	Asn Ile Ala Thr	Tyr	Tyr Gly Ala Phe	Ile
	80		85		90
Lys Lys Ser Pro	Pro	Gly His Asp Asp	Gln	Leu Trp Leu Val	Met
	95		100		105
Glu Phe Cys Gly	Ala	Gly Ser Ile Thr	Asp	Leu Val Lys Asn	Thr
	110		115		120
Lys Gly Asn Thr	Leu	Lys Glu Asp Trp	Ile	Ala Tyr Ile Ser	Arg
	125		130		135
Glu Ile Leu Arg	Gly	Leu Ala His Leu	His	Ile His His Val	Ile
	140		145		150
His Arg Asp Ile	Lys	Gly Gln Asn Val	Leu	Leu Thr Glu Asn	Ala

	155		160		165
Glu Val Lys Leu	Val Asp Phe Gly Val	Ser Ala Gln Leu Asp Arg			
	170		175		180
Thr Val Gly Arg	Arg Asn Thr Phe Ile	Gly Thr Pro Tyr Trp Met			
	185		190		195
Ala Pro Glu Val	Ile Ala Cys Asp Glu	Asn Pro Asp Ala Thr Tyr			
	200		205		210
Asp Tyr Arg Ser	Asp Leu Trp Ser Cys	Gly Ile Thr Ala Ile Glu			
	215		220		225
Met Ala Glu Gly	Ala Pro Pro Leu Cys	Asp Met His Pro Met Arg			
	230		235		240
Ala Leu Phe Leu	Ile Pro Arg Asn Pro	Pro Pro Arg Leu Lys Ser			
	245		250		255
Lys Lys Trp Ser	Lys Lys Phe Phe Ser	Phe Ile Glu Gly Cys Leu			
	260		265		270
Val Lys Asn Tyr	Met Gln Arg Pro Ser	Thr Glu Gln Leu Leu Lys			
	275		280		285
His Pro Phe Ile	Arg Asp Gln Pro Asn	Glu Arg Gln Val Arg Ile			
	290		295		300
Gln Leu Lys Asp	His Ile Asp Arg Thr	Arg Lys Lys Arg Gly Glu			
	305		310		315
Lys Asp Glu Thr	Glu Tyr Glu Tyr Ser	Gly Ser Glu Glu Glu Glu			
	320		325		330
Glu Glu Val Pro	Glu Gln Glu Gly Glu	Pro Ser Ser Ile Val Asn			
	335		340		345
Val Pro Gly Glu	Ser Thr Leu Arg Arg	Asp Phe Leu Arg Leu Gln			
	350		355		360
Gln Glu Asn Lys	Glu Arg Ser Glu Ala	Leu Arg Arg Gln Gln Leu			
	365		370		375
Leu Gln Glu Gln	Gln Leu Arg Glu Gln	Glu Glu Tyr Lys Arg Gln			
	380		385		390
Leu Leu Ala Glu	Arg Gln Lys Arg Ile	Glu Gln Gln Lys Glu Gln			
	395		400		405
Arg Arg Arg Leu	Glu Glu Gln Gln Arg	Arg Glu Arg Glu Ala Arg			
	410		415		420
Arg Gln Gln Glu	Arg Glu Gln Arg Arg	Arg Glu Gln Glu Glu Lys			
	425		430		435
Arg Arg Leu Glu	Glu Leu Glu Arg Arg	Arg Lys Glu Glu Glu Glu			
	440		445		450
Arg Arg Arg Ala	Glu Glu Glu Lys Arg	Arg Val Glu Arg Glu Gln			
	455		460		465
Glu Tyr Ile Arg	Arg Gln Leu Glu Glu	Glu Gln Arg His Leu Glu			
	470		475		480
Val Leu Gln Gln	Gln Leu Leu Gln Glu	Gln Ala Met Leu Leu His			
	485		490		495
Asp His Arg Arg	Pro His Pro Gln His	Ser Gln Gln Pro Pro Pro			
	500		505		510
Pro Gln Gln Glu	Arg Ser Lys Pro Ser	Phe His Ala Pro Glu Pro			
	515		520		525
Lys Ala His Tyr	Glu Pro Ala Asp Arg	Ala Arg Glu Val Glu Asp			
	530		535		540
Arg Phe Arg Lys	Thr Asn His Ser Ser	Pro Glu Ala Gln Ser Lys			
	545		550		555
Gln Thr Gly Arg	Val Leu Glu Pro Pro	Val Pro Ser Arg Ser Glu			
	560		565		570
Ser Phe Ser Asn	Gly Asn Ser Glu Ser	Val His Pro Ala Leu Gln			
	575		580		585
Arg Pro Ala Glu	Pro Gln Val Pro Val	Arg Thr Thr Ser Arg Ser			
	590		595		600
Pro Val Leu Ser	Arg Arg Asp Ser Pro	Leu Gln Gly Ser Gly Gln			
	605		610		615
Gln Asn Ser Gln	Ala Gly Gln Arg Asn	Ser Thr Ser Ser Ile Glu			
	620		625		630
Pro Arg Leu Leu	Trp Glu Arg Val Glu	Lys Leu Val Pro Arg Pro			
	635		640		645
Gly Ser Gly Ser	Ser Ser Gly Ser Ser	Asn Ser Gly Ser Gln Pro			
	650		655		660

Gly Ser His Pro	Gly Ser Gln Ser Gly	Ser Gly Glu Arg Phe	Arg
665	670	675	
Val Arg Ser Ser	Ser Lys Ser Glu Gly	Ser Pro Ser Gln Arg	Leu
680	685	690	
Glu Asn Ala Val	Lys Lys Pro Glu Asp	Lys Lys Glu Val Phe	Arg
695	700	705	
Pro Leu Lys Pro	Ala Gly Glu Val Asp	Leu Thr Ala Leu Ala	Lys
710	715	720	
Glu Leu Arg Ala	Val Glu Asp Val Arg	Pro Pro His Lys Val	Thr
725	730	735	
Asp Tyr Ser Ser	Ser Ser Glu Glu Ser	Gly Thr Thr Asp Glu	Glu
740	745	750	
Asp Asp Asp Val	Glu Gln Glu Gly Ala	Asp Glu Ser Thr Ser	Gly
755	760	765	
Pro Glu Asp Thr	Arg Ala Ala Ser Ser	Leu Asn Leu Ser Asn	Gly
770	775	780	
Glu Thr Glu Ser	Val Lys Thr Met Ile	Val His Asp Asp Val	Glu
785	790	795	
Ser Glu Pro Ala	Met Thr Pro Ser Lys	Glu Gly Thr Leu Ile	Val
800	805	810	
Arg Gln Thr Gln	Ser Ala Ser Ser Thr	Leu Gln Lys His Lys	Ser
815	820	825	
Ser Ser Ser Phe	Thr Pro Phe Ile Asp	Pro Arg Leu Leu Gln	Ile
830	835	840	
Ser Pro Ser Ser	Gly Thr Thr Val Thr	Ser Val Val Gly Phe	Ser
845	850	855	
Cys Asp Gly Met	Arg Pro Glu Ala Ile	Arg Gln Asp Pro Thr	Arg
860	865	870	
Lys Gly Ser Val	Val Asn Val Asn Pro	Thr Asn Thr Arg Pro	Gln
875	880	885	
Ser Asp Thr Pro	Glu Ile Arg Lys Tyr	Lys Lys Arg Phe Asn	Ser
890	895	900	
Glu Ile Leu Cys	Ala Ala Leu Trp Gly	Val Asn Leu Leu Val	Gly
905	910	915	
Thr Glu Ser Gly	Leu Met Leu Leu Asp	Arg Ser Gly Gln Gly	Lys
920	925	930	
Val Tyr Pro Leu	Ile Asn Arg Arg Arg	Phe Gln Gln Met Asp	Val
935	940	945	
Leu Glu Gly Leu	Asn Val Leu Val Thr	Ile Ser Gly Lys Lys	Asp
950	955	960	
Lys Leu Arg Val	Tyr Tyr Leu Ser Trp	Leu Arg Asn Lys Ile	Leu
965	970	975	
His Asn Asp Pro	Glu Val Glu Lys Lys	Gln Gly Trp Thr Thr	Val
980	985	990	
Gly Asp Leu Glu	Gly Cys Val His Tyr	Lys Val Val Lys Tyr	Glu
995	1000	1005	
Arg Ile Lys Phe	Leu Val Ile Ala Leu	Lys Ser Ser Val Glu	Val
1010	1015	1020	
Tyr Ala Trp Ala	Pro Lys Pro Tyr His	Lys Phe Met Ala Phe	Lys
1025	1030	1035	
Ser Phe Gly Glu	Leu Val His Lys Pro	Leu Leu Val Asp Leu	Thr
1040	1045	1050	
Val Glu Glu Gly	Gln Arg Leu Lys Val	Ile Tyr Gly Ser Cys	Ala
1055	1060	1065	
Gly Phe His Ala	Val Asp Val Asp Ser	Gly Ser Val Tyr Asp	Ile
1070	1075	1080	
Tyr Leu Pro Thr	His Ile Gln Cys Ser	Ile Lys Pro His Ala	Ile
1085	1090	1095	
Ile Ile Leu Pro	Asn Thr Asp Gly Met	Glu Leu Leu Val Cys	Tyr
1100	1105	1110	
Glu Asp Glu Gly	Val Tyr Val Asn Thr	Tyr Gly Arg Ile Thr	Lys
1115	1120	1125	
Asp Val Val Leu	Gln Trp Gly Glu Met	Pro Thr Ser Val Ala	Tyr
1130	1135	1140	
Ile Arg Ser Asn	Gln Thr Met Gly Trp	Gly Glu Lys Ala Ile	Glu
1145	1150	1155	
Ile Arg Ser Val	Glu Thr Gly His Leu	Asp Gly Val Phe Met	His

	1160		1165		1170
Lys Arg Ala Gln Arg	Leu Lys Phe Leu Cys	Glu Arg Asn Asp Lys			
	1175		1180		1185
Val Phe Phe Ala Ser	Val Arg Ser Gly Gly	Ser Ser Gln Val Tyr			
	1190		1195		1200
Phe Met Thr Leu Gly	Arg Thr Ser Leu Leu	Ser Trp			
	1205		1210		

<210> 10
 <211> 280
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5046791CD1

<400> 10	
Met Gln Pro Leu Arg	Val Asn Ser Gln Pro Gly Pro Gln Lys Arg
1	5
Cys Leu Phe Val Cys	Arg His Gly Glu Arg Met Asp Val Val Phe
	20
Gly Lys Tyr Trp Leu	Ser Gln Cys Phe Asp Ala Lys Gly Arg Tyr
	35
Ile Arg Thr Asn Leu	Asn Met Pro His Ser Leu Pro Gln Arg Ser
	50
Gly Gly Phe Arg Asp	Tyr Glu Lys Asp Ala Pro Ile Thr Val Phe
	65
Gly Cys Met Gln Ala	Arg Leu Val Gly Glu Ala Leu Leu Glu Ser
	80
Asn Thr Ile Ile Asp	His Val Tyr Cys Ser Pro Ser Leu Arg Cys
	95
Val Gln Thr Ala His	Asn Ile Leu Lys Gly Leu Gln Gln Glu Asn
	110
His Leu Lys Ile Arg	Val Glu Pro Gly Leu Phe Glu Trp Thr Lys
	125
Trp Val Ala Gly Ser	Thr Leu Pro Ala Trp Ile Pro Pro Ser Glu
	140
Leu Ala Ala Ala Asn	Leu Ser Val Asp Thr Thr Tyr Arg Pro His
	155
Ile Pro Ile Ser Lys	Leu Val Val Ser Glu Ser Tyr Asp Thr Tyr
	170
Ile Ser Arg Ser Phe	Gln Val Thr Lys Glu Ile Ile Ser Glu Cys
	185
Lys Ser Lys Gly Asn	Asn Ile Leu Ile Val Ala His Ala Ser Ser
	200
Leu Glu Ala Cys Thr	Cys Gln Leu Gln Gly Leu Ser Pro Gln Asn
	215
Ser Lys Asp Phe Val	Gln Met Val Arg Lys Ile Pro Tyr Leu Gly
	230
Phe Cys Ser Cys Glu	Glu Leu Gly Glu Thr Gly Ile Trp Gln Leu
	245
Thr Asp Pro Pro Ile	Leu Pro Leu Thr His Gly Pro Thr Gly Gly
	260
Phe Asn Trp Arg Glu	Thr Leu Leu Gln Glu
	275

<210> 11
 <211> 114
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1416174CD1

<400> 11

```

Met Leu Ala Ile Ser Pro Ser His Leu Gly Ala Asp Leu Val Ala
1      5      10      15
Ala Pro His Ala Arg Phe Asp Asp Gly Leu Val His Leu Cys Trp
20     25     30
Val Arg Thr Gly Ile Ser Arg Ala Ala Leu Leu Arg Leu Phe Leu
35     40     45
Ala Met Glu Arg Gly Ser His Phe Ser Leu Gly Cys Pro Gln Leu
50     55     60
Gly Tyr Ala Ala Ala Arg Ala Phe Arg Leu Glu Pro Leu Thr Pro
65     70     75
Arg Gly Val Leu Thr Val Asp Gly Glu Gln Val Glu Tyr Gly Pro
80     85     90
Leu Gln Ala Gln Met His Pro Gly Ile Gly Thr Leu Leu Thr Gly
95     100    105
Pro Pro Gly Cys Pro Gly Arg Glu Pro
110

```

<210> 12
 <211> 375
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3244919CD1

```

<400> 12
Met Gly Ser Ser Met Ser Ala Ala Thr Ala Arg Arg Pro Val Phe
1      5      10      15
Asp Asp Lys Glu Asp Val Asn Phe Asp His Phe Gln Ile Leu Arg
20     25     30
Ala Ile Gly Lys Gly Ser Phe Gly Lys Val Cys Ile Val Gln Lys
35     40     45
Arg Asp Thr Glu Lys Met Tyr Ala Met Lys Tyr Met Asn Lys Gln
50     55     60
Gln Cys Ile Glu Arg Asp Glu Val Arg Asn Val Phe Arg Glu Leu
65     70     75
Glu Ile Leu Gln Glu Ile Glu His Val Phe Leu Val Asn Leu Trp
80     85     90
Tyr Ser Phe Gln Asp Glu Glu Asp Met Phe Met Val Val Asp Leu
95     100    105
Leu Leu Gly Gly Asp Leu Arg Tyr His Leu Gln Gln Asn Val Gln
110    115    120
Phe Ser Glu Asp Thr Val Arg Leu Tyr Ile Cys Glu Met Ala Leu
125    130    135
Ala Leu Asp Tyr Leu Arg Gly Gln His Ile Ile His Arg Asp Val
140    145    150
Lys Pro Asp Asn Ile Leu Leu Asp Glu Arg Gly His Ala His Leu
155    160    165
Thr Asp Phe Asn Ile Ala Thr Ile Ile Lys Asp Gly Glu Arg Ala
170    175    180
Thr Ala Leu Ala Gly Thr Lys Pro Tyr Met Ala Pro Glu Ile Phe
185    190    195
His Ser Phe Val Asn Gly Gly Thr Gly Tyr Ser Phe Glu Val Asp
200    205    210
Trp Trp Ser Val Gly Val Met Ala Tyr Glu Leu Leu Arg Gly Trp
215    220    225
Arg Pro Tyr Asp Ile His Ser Ser Asn Ala Val Glu Ser Leu Val
230    235    240
Gln Leu Phe Ser Thr Val Ser Val Gln Tyr Val Pro Thr Trp Ser
245    250    255
Lys Glu Met Val Ala Leu Leu Arg Lys Leu Leu Thr Val Asn Pro
260    265    270
Glu His Arg Leu Ser Ser Leu Gln Asp Val Gln Ala Ala Pro Ala
275    280    285
Leu Ala Gly Val Leu Trp Asp His Leu Ser Glu Lys Arg Val Glu
290    295    300

```

Pro	Gly	Phe	Val	Pro	Asn	Lys	Gly	Arg	Leu	His	Cys	Asp	Pro	Thr
				305					310					315
Phe	Glu	Leu	Glu	Glu	Met	Ile	Leu	Glu	Ser	Arg	Pro	Leu	His	Lys
				320					325					330
Lys	Lys	Lys	Arg	Leu	Ala	Lys	Asn	Lys	Ser	Arg	Asp	Asn	Ser	Arg
				335					340					345
Asp	Ser	Ser	Gln	Ser	Ala	Pro	Arg	Ser	Lys	Ser	Lys	Pro	Ser	Thr
				350					355					360
Gln	Arg	Gln	Gly	Ser	Trp	Ala	Leu	Ala	Ser	Ser	Gly	Leu	Gly	Glu
				365					370					375

<210> 13
 <211> 1859
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 058860CB1

<400> 13
 gagagataact ccacaccccc aggagagact ctagagagat attccacacc cccaggagag 60
 actctggaga gatactccac acccccagga gagactctag agcgatattc cacaccccca 120
 ggggaggcac tagagagata ttctattcct actggaggac caaacccac tggtaactttt 180
 aaaacatata catcaaaaat agaaatggaa gacggtagac caaatgagca tttctacaca 240
 cctacagaag agaggggttc agcttatgaa atatggcgtt ccgattcatt tggtagacccc 300
 aatgaagcca ttgagccaaa agacaatgaa atgcctccat cttttattga acctctgacc 360
 aaaaggaagg tatatgaaaa cacaacacta ggcttcattg ttgaagttga aggtcttcca 420
 gttcctgggtg tgaaatggta tcgaaataaa tctttactag agccagatga aagaatcaaa 480
 atggaaagag tgggtaattgt gtgttcaact gaaatttcta acattcaaaa aggagaaggg 540
 ggagagtaca tgtgtcatgc tgtaaacatc ataggggaag caaagagctt tgcaaatgta 600
 gacataatgc cccaggaaga aagagtgggt gcactaccac ctccagtaac acatcagcat 660
 gtcatggagt ttgatttgga acacaccaca tcatcaagaa caccttctcc tcaagaaatt 720
 gtcttggaag ttgaattaag tgaaaagac gttaaagaat ttgagaagca ggtgaaaata 780
 gtgacagttc ccgaatttac tcttgaccat aaaagtatga ttgtgagtct agatgttctt 840
 ccatttaatt ttgtagatcc aaatatggat tcaagggagg gagaagacaa agaactaaaa 900
 attgatttag aagtatttga aatgcctcct cgctttataa tgcctatttg tgattttaaa 960
 attccagaaa attcagatgc tgtattcaaa tgttcagtca tagggatccc gactcccga 1020
 gttaagtggg ataaagaata tatgtgtatt gagccagata atattaaata cgtgatttag 1080
 gaggagaagg gaagtcacac tcttaaaatt cgaaatgtct gtctttctga tagtgcaaca 1140
 tacaggtgca gagctgtgaa ttgtgtagga gaggtatct gtctggggatt cctcaccatg 1200
 ggagattctg aaatatttgc tgtgatagca aagaaaagca aagtgaactt aagcagttta 1260
 atggaagaat tggctctaaa gagcaactac acagacagtt tttttgaatt tcagggtgggt 1320
 gaagggcctc ccaggtttat caaagggtatt tctgactgtt atgcaccaat aggtacagca 1380
 gcatattttc agtgcttagt tctgtgctct ccaagaccca cggtttactg gtacaaagat 1440
 ggaaaattag tccaaggaag aaggttcaact gttgaggaaa gtggcacagg gttccataac 1500
 ctgtttataa caagcttagt aaagagtgtg gaaggagagt ataggtgtgt agctacaaac 1560
 aaatcaggaa tggctgagag ctttgcagca ctacacctaa cttaaaatgt aatgttttag 1620
 tgcctcagta attattagca ttgatctgag tgctttcata ttttccaaat tatgtggatc 1680
 taataaactt ccaaacaggt ccaccatatt tgaattcatt accttgagga cccctaaaga 1740
 aataatctct atgtagaaat ctcactcttg taatacatgt aaatattttg ttatctgaac 1800
 tgtggaatca tcacttgtgt caatcatgct gtgtaatatc aaacacaatt aaatctctc 1859

<210> 14
 <211> 3501
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2041716CB1

<400> 14
 gtggtgtggc tgcagtggag agttcccaac aaggctacgc agaagaaccc ccttgactga 60
 agcaatggag gggggtccag ctgtctgctg ccaggatcct cgggcagagc tggtagaacg 120
 ggtggcagcc atcgatgtga ctcacttggg ggaggcagat ggtggcccag agcctactag 180

aaacggtgtg gacccccac cacgggccag agctgcctct gtgatccctg g
 aagactgtct ccagccggc ctagcctctc agccaggaag ctttccctac agy/35304
 agcaggaagc tatctggagg cgcaggctgg gccttatgcc acggggcctg cca
 ctcccccg gcctggcgga ggcccacat ctagtccac cacgtggcca tctca
 agaggactgc gtgcagctga accagtacaa gctgcagagt gagattggca aggtggg
 gactgatgcc tatctgcagg gtgcttacgg tgtggtgagg ctggcctaca acgaaagt
 agacagacac tatgcaatga aagtcctttc caaaaagaag ttactgaagc agtatggct
 tccacgtcgc cctccccga gaggggtccca ggctgcccag ggaggaccag ccaagcagct
 gctgcccctg gagcgggtgt accaggagat tgccatcctg aagaagctgg accacgtgaa 7
 tgtggtcaaa ctgatcgagg tcctggatga cccagctgag gacaacctct atttgggtga 780
 cctcctgaga aaggggcccg tcctggaagt gccctgtgac aagccttct cggaggagca 840
 agctgcctc tacctgcggg acgtcatcct gggcctcgag tacttgact gccagaagat 900
 cgtccacagg gacatcaagc catccaacct gctcctgggg gatgatggg cagcacggc 960
 cgccgacttt ggcgtcagca accagtttga ggggaacgac gctcagctgt ccagcaggg 1020
 gggaaccca gattcatgg cccccaggc catttctgat tccggccaga gcttcagtgg 1080
 gaaggccttg gatgtatgg ccactggcgt cacgttgtac tgctttgtct atgggaagt 1140
 cccgttcatc gacgatttca tcctggcct ccacaggaag atcaagaatg agcccggtgg 1200
 gtttctctgag gagccagaaa tcagcgagga gctcaaggac ctgatcctga agatgttaga 1260
 caagaatccc gagacgagaa ttgggggtgcc agacatcaag ttgcacctt gggtgaccaa 1320
 gaacggggag gagccccctt cttcgaggga ggagcactgc agcgtgggtg aggtgacaga 1380
 ggaggaggtt aagaactcag tcaggctcat cccagctgg accacgggtg tcctgggtgaa 1440
 gtccatgctg aggaagcgtt cctttgggaa cccgtttgag cccaagcac ggagggaaga 1500
 gcatccatg tctgctccag gaaacctact ggtgaaagaa ggggttgggt aagggggcaa 1560
 gagccagag ctccccggcg tccagggaaga cgaggctgca tcttgagccc ctgcatgcac 1620
 ccagggccac cgggcagcac actcatcccg gcagccccac ggcccacccc tcatgcaaca 1680
 gccgcccccg caggcagggg gctggggact gcagccccac tcccgcctt ccccatcgt 1740
 gctgcatgac ctccacgac gcacgtccag ggacagactg gaatgtatgt catttgggt 1800
 cttgggggca gggctccac gaggccatc tcctcttctt ggacctcctt ggcttgacc 1860
 attctgtggg gaaaccgggt gcccatggag cctcagaaat gccaccggc tggttggcat 1920
 ggcctggggc agggagcaga ggcaggagac caagatggca ggtggaggcc aggttacca 1980
 caacggaaga gacctccgc tggggccggg caggcctggc tcagctgcca caggcatatg 2040
 gtggagaggg ggggtacctg cccaccttgg ggtggtggca ccagagctct tgtctattca 2100
 gacgttggtg tgggggctcg gacccctcac tggggacagg gccagtgtt gagaatctg 2160
 attccttttt tgttgtctt tacttttgtt tttaacctgg ggggtcgggg agaggccctg 2220
 cttgggaaca tctcacgagc tttctacat cttcgtgggt tcccagcaca gccaagatt 2280
 atttggcagc caagtggatg gaactaact tcctggactg tgtttcgcat tcggcggtat 2340
 ctggaagtgt gactgaacgg aatcaagctc tgagcagagg cctgaagcgg aagcaccaca 2400
 tctgccccgc ccactctact ctctccctg atgatgcccc tagagctgag gctggagaag 2460
 acaccaggc tgactttgac cgagggccat ggacgcgaca ggcctgtggc cctgcgcatg 2520
 ctgaaataac tggaaaccag cctctcctcc tacaccggc taccatctg ggcccagag 2580
 ctgcactcac actcctacaa cgaaggacaa actgtccagg tggagggat cagcagacac 2640
 agaacttga ggggtgtgca cgtggcagg tgccctctgc ggcaattgcc tcaccctgag 2700
 gacatcagca gtcagcctgc tcagagcggg ggtgctggag cgcgtgcaga cacagctctt 2760
 ccggagcagc cttcacctt cctctgggat cagtgtccgg ctggccgacg tggcatttgc 2820
 tgaccgaatg ctcatagagg ttgacccca cagggtcacg caggactcgg acactgcc 2880
 ggaaacatgg atggacaagg gcttttggcc acaggtgtgg gtgtcctgtt ggaggaggc 2940
 ttgttttgag aaggagggtt ggcctgggga gaaaccgga tcccgctgca tctccgccc 3000
 tgtgggtgca tgtcgctgc tcactgttg cacacagctc actcgtatgt cctgactgg 3060
 tacatgcatc tgaataacag tttctacgtc tatttaaggc taggagccga atgtgcccc 3120
 ttgtcagtgg gtccacgttt ctccccggct cctctgggct agggcagtgt ggcccaggc 3180
 ttaaaaagtt actcgtact gtttttaaga acacttttat agagttagtg gaaggcaagt 3240
 taagagccaa tcactgatcc ccaagtgtt cttgagcatc tggctctggg ggaccacttt 3300
 gatcggacc acccttgga agctcaggg taggccagg tgggatgctc accctgtcac 3360
 tgagggtttt ggttgcatc gttgtttttg aatgtagcac aagcgatgag caaactctat 3420
 aagagtgttt taaaaattaa cttcccagga agtgagttaa aaacaataaa agccctttct 3480
 tgagttaaaa agaaaaaaa a

<210> 15
 <211> 3039
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7472005CB1
 <400> 15


```

atggcccccg cccgggggccc cctgccccct gcgctctggg tcgtcacggc cgcgggggcg 60
gcggccacct gcgtgtccgc ggcgcgcgcc gaagtgaatt tgctggacac gtcgaccatc 120
cacgggggact ggggctggct cacgtatccg gctcatgggt gggactccat caacgaggtg 180
gacgagtcct tccagcccat ccacacgtac caggtttgca acgtcatgag ccccaaccag 240
aacaactggc tgcgcacgag ctgggtcccc cgagacggcg cccggcgcg tctatgctgag 300
atcaagttta cccctgcgca ctgcaacagc atgcctgggt tgctgggcac ctgcaaggag 360
accttcaacc tctactacct ggagtcggac cgcgacctgg gggccagcac acaagaaagc 420
cagttcctca aaatcgacac cattgcgggc gacgagagct tcacagggtg cgaccttggg 480
gtgcggcgct tcaagctcaa cacggaggtg cgcagtgtgg gtccctcag caagcgcgcg 540
ttctacctgg ccttccagga cataggtgcc tgctggcca tctctctct cgcctctac 600
tataagaagt gccctgccat ggtgcgcaat ctggctgcct tctcggaggg agtgacgggg 660
gccgactcgt cctcactggt ggaggtgagg ggccagtgcg tgcggcactc agaggagcgg 720
gacacaccca agatgtactg cagcgcgagg ggcgagtggc tcgtgcccat cggcaaatgc 780
gtgtgcagtg cgggtacga ggagcggcgg gatgcctgtg tggcctgtga gctgggcttc 840
tacaagtcag cccctgggga ccagctgtgt gcccgctgcc ctccccacag cactccgca 900
gtccagccg cccaagcctg ccactgtgac ctacgtact accgtgcagc cctggaccgg 960
ccgtcctcag cctgcaccgg gccaccctcg gcaccagtga acctgatctc cagtgtgaat 1020
gggacatcag tgactctgga gtggggccct cccctggacc cagggtggcg cagtgcacac 1080
acctacaatg ccgtgtgccg ccgctgcccc tgggcactga gccgctgcca ggcatgtggg 1140
agcggcacc gctttgtgcc ccagcagaca agcctggtgc aggcagcct gctgggtggc 1200
aacctgtcgg ccccatgaa ctactccttc tggatcgagg ccgtcaatgg cgtgtccgag 1260
ctgagccccg agccccgccc ggccgctgtg gtcaacatca ccacgaacca ggcagccccg 1320
tcccaggtgg tgggtgatccg tcaagagcgg gcggggcaga ccagcgtctc gctgctgtgg 1380
caggagcccc agcagccgaa cggcatcatc ctggagtatg agatcaagta ctacgagaag 1440
gacaaggaga tgcagagcta ctccaccctc aaggccgtca ccaccagagc caccgtctcc 1500
ggcctcaagc cgggcaccgg ctactgttcc caggctcgag cccgcacctc agcaggctgt 1560
ggccgcttca gccaggccat ggaggtggag accgggaaac cccggccccg ctatgacacc 1620
aggaccattg tctggatctg cctgacgctc atcacgggcc tgggtgtgct tctgctcctg 1680
ctcatctgca agaagaggca ctgtggctac agcaaggcct tccaggactc ggacgaggag 1740
aagatgcact atcagaatgg acaggcacc ccacctgtct tctgcctct gcacacccc 1800
ccgggaaagc tcccagagcc ccagttctat gcggaacccc acacctacga ggagccaggc 1860
cgggcggggc gcagtttcac tcgggagatc gaggcctcta ggatccacat cgagaaaatc 1920
atcggtctct gagactccgg ggaagtctgc tacgggaggc tgcggtgccc agggcagcgg 1980
gatgtgcccc tggccatcaa ggccctcaaa gccggctaca cggagagaca gaggcgggac 2040
ttctgagcgg aggcgtccat catggggcaa ttcgaccatc ccaacatcat ccgctcagag 2100
ggtgtcgtca cccgtggccg cctggcaatg attgtgactg agtacatgga gaacggctct 2160
ctggacacct tctgaggag ccacgacggg cagttcacca tcatgcagct ggtgggcatg 2220
ctgagaggag tgggtgccgg catgcgtac ctctcagacc tgggctatgt ccaccgagac 2280
ctggccgccc gcaacgtcct ggttgacagc aacctggtct gcaagggtgc tgacttcggg 2340
ctctcacggg tgctggagga cgaccgggat gctgcctaca ccaccacggg cgggaagatc 2400
cccatccgct ggacggcccc agaggccatc gccttcgca ccttctcctc ggccagcgac 2460
gtgtggagct tcggcggtgt catgtgggag gtgctggcct atggggagcg gccctactgg 2520
aacatgacca accgggatgt gagtgcgaag ccctggcagg tcatcagctc tgtggaggag 2580
gggtaccgcc tgcccgccc catgggtgc cccacgccc tgcaccagct catgctcgac 2640
tggtggcaca aggacggggc gcagcggcct cgcttctccc agattgtcag tgtcctcgat 2700
gcgtcatccc gcagccctga gagtctcagg gccaccgcca cagtcagcag gtgcccaccc 2760
cctgccttcg tccggagctg ctttgacctc cgagggggca gcggtggcgg tgggggcctc 2820
accgtggggg actggctgga ctccatccgc atgggcgggt accgagacca cttcgctgag 2880
ggcgataact cctctctggg catggtgcta cgcatgaacg cccaggacgt gcgcgccctg 2940
ggcatcacc tcatgggcca ccagaagaag atcctgggca gcattcagac catgcggggc 3000
cagctgacca gcacccaggg gccccggcg cacctctga 3039

```

<210> 16

<211> 1104

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472006CB1

<400> 16

```

atggatgacg ctgctgtcct caagcgacga ggctacctcc tggggataaa tttaggagag 60
ggctcctatg caaaagtaaa atctgcttac tctgagcgcc tgaagttcaa tgtggcgatc 120
aagatcatcg accgcaagaa ggccccgca gacttcttgg agaaattcct tccccgggaa 180
attgagattc tggccatgtt aaaccactgc tccatcatta agacctacga gatctttgag 240
acatcacatg gcaaggctta catcgtcatg gagctcgag tccaggcgga cctcctcgag 300

```

ttaatcaaaa	cccggggagc	cctgcatgag	gacgaagctc	gcaagaagtt	ccaccagctt	360
tccttgccca	tcaagtactg	ccacgacctg	gacgtcgtcc	accgggacct	caagtgtgac	420
aaccttctcc	ttgacaagga	cttcaacatc	aagctgtccg	acttcagctt	ctccaagcgc	480
tgcttgccgg	atgacagtgg	tcgaatggcc	ttaagcaaga	ccttctgtgg	gtcaccagcg	540
tatcgccccc	cagaggtgct	gcagggcatt	ccctaccagc	ccaaggtgta	cgacatctgg	600
agcctaggcg	tgatcctcta	catcatggtc	tgccgtctca	tgccctacga	cgactccaac	660
atcaagaaga	tgctgcgtat	ccagaaggag	caccgcgtca	acttcccacg	ctccaagcac	720
ctgacaggcg	agtgaaggga	cctcatctac	cacatgctgc	agcccagcgt	caaccggcgg	780
ctccacatcg	acgagatcct	cagccactgc	tggatgcagc	ccaaggcacg	gggatctccc	840
tctgtggcca	tcaacaagga	gggggagagt	tcccggggaa	ctgaaccctt	gtggaccccc	900
gaacctggct	ctgacaagaa	gtctgccacc	aagctggagc	ctgagggaga	ggcacagccc	960
caggcacagc	ctgagacaaa	acccgagggg	acagcaatgc	aaatgtccag	gcagtccggag	1020
atcctgggtt	tccccagcaa	gccgtcgact	atggagacag	aggaaggggc	cccccaacag	1080
cctccagaga	cgccggccca	gtga				1104

<210> 17

<211> 3939

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2902460CB1

<400> 17

ccgcagtgtg	ctggaaaggc	agctgcggca	gtagcgtgag	cagcccaagt	tgggctggtc	60
gcctgcgagg	ggaccggcag	caggtgggtg	cagccgggtac	cctctccccg	ccaggccgga	120
ggaggccaag	aggaagctgc	ggatcttgca	gcgcgagttg	cagaacgtgc	aggtgaacca	180
gaaagtgggc	atgtttgagg	cgacatcca	ggcacagagc	tccgccattc	aagcgccccg	240
cagcccgctg	ttgggcaggg	ctcgctcgcc	ctccccgtgc	cccttccgca	gcagcagtca	300
gccccctgga	agggctcctg	ttcagggcgc	ccggagcgag	gaacggaggga	caaagtccctg	360
gggggagcaa	tgtccagaga	cttcaggaaac	cgactccggg	aggaaggagg	ggcccagcct	420
atgctcctcg	caggtgaaga	aaggaaatgcc	acctcttccc	ggccgggctg	cccctacagg	480
atcagaggct	cagggctccat	ccgcttttgt	aaggatggag	aagggtatcc	ctgccagtcc	540
ccgctgtggc	tcacccacag	ctatggaaat	tgacaaaagg	ggctctccta	ccccgggaac	600
tcggagctgc	ctagctccct	cattggggct	gttcggagct	agcttaacga	tggccacgga	660
agtggcagcg	aggaattacat	ccactgggcc	acacgcgtca	caggatcttg	ccctcactga	720
gccgtctggg	agagcccgtg	agcttgaggga	cctgcagccc	ccagaggccc	tgggtggagag	780
gcaggggcag	tttctgggca	gtgagacaag	cccagcccca	gaaaggggcg	ggccccgcga	840
tggagaaccc	cctgggaaga	tggggaaagg	atatctgccc	tgtggcatgc	cgggctctgg	900
ggagcttgaa	gtgggcaaaa	ggccagagga	gacgactgtg	agcgtgcaaa	gcgcagagtc	960
ctctgatgcc	ctgagctggg	ccaggctgcc	cagggccctg	gcctccgtag	gccctgagga	1020
ggccccgaag	ggggcccccg	tgggcggggg	gcgttggcag	ctctccgaca	gagtggaggg	1080
aggggtcccca	acgctgggct	tgcctggggg	cagcccttca	gcacagccgg	ggaccgggaa	1140
tgtggagcgg	ggaattcctt	ctggcagaat	gctggagcct	tgccctgtt	gggacgctgc	1200
gaaagatctg	aaagaacctc	agtgccctcc	tggggacagg	gtgggtgtgc	agcctgggaa	1260
ctccaggggt	tggcagggca	ccatggagaa	agccgggttg	gcttggacgc	gtggcacagg	1320
ggtgcaatca	gaggggactt	gggaaagcca	gcggcaggac	agtgatgccc	tcccaagtc	1380
ggagctgcta	ccccaaagtc	aggacaagcc	tttcttgagg	aaggcctgca	gccccagcaa	1440
catacctgct	gtcatcatta	cagacatggg	cacccaggag	gatggggcct	tggaggagac	1500
gcagggaagc	cctcggggca	acctgcccct	gaggaactg	tcctcttctc	cggcctctc	1560
cacgggcttc	tcctcactct	acgaagactc	agaggaggac	atctccagt	accctgagcg	1620
caccttggtc	cccaactcag	ccttcttgca	tacctggac	cagcagaaac	ctagagtga	1680
caaatcatgg	aggaagataa	aaaacatggt	gcactggtct	cccttcgtca	tgtccttcaa	1740
gaagaagtac	ccctggatcc	agctggcagg	acacgcaggg	agtttcaagg	cagctgccaa	1800
tggcaggatc	ctgaagaagc	actgtgagtc	agagcagcgc	tgccctggacc	ggctgatggg	1860
ggatgtgctg	aggcccttcg	tacctgccta	ccatggggat	gtgggtgaagg	acggggagcg	1920
ctacaaccag	atggacgacc	tgttgccgca	cttcgactcg	ccctgtgtga	tggactgcaa	1980
gatgggaaatc	aggacctacc	tggaggagga	gctcacaag	gcccgggaaga	agccagcct	2040
gcgggaaggac	atgtaccaga	agatgatcga	ggtggacccc	gaggccccca	ccgaggagga	2100
aaaagcacag	cgggctgtga	ccaagccacg	gtacatgcag	tggcggggaga	ccatcagctc	2160
cacggccacc	ctggggttca	ggatcgaggg	aatcaagaaa	gaagacggca	ccgtgaaccg	2220
ggacttcaag	aagacaaaaa	cgaggggagca	ggtcaccgag	gccttcagag	agttcactaa	2280
aggaaccatc	aacatcctga	tcgcctatcg	ggaccggctg	aaggccattc	gaaccactct	2340
agaagtttct	cccttcttca	agtgccacga	ggtcattggc	agctccctcc	tcttcattca	2400
cgacaagaag	gaacaggcca	aagtgtggat	gatcgacttt	gggaaaacca	cggccctgcc	2460
tgaggggccag	accctgcagc	atgacgtccc	ctggcaggag	gggaaccggg	aggatggcta	2520

```

cctctcgggg ctcaataacc tcgtcgacat cctgaccgag atgtcccagg atgccccact 2580
cgccctgagct gcccacgccc tccttggecc ccgcctgggc ctcttttcc cctcctgtgc 2640
ttcctttctc gttcctaact ttctcttcac ttacacctga ctgacctcc tgaactgcac 2700
tacaagacac tttgtagaag aggagatgag agttttctagt cattttccta acttcagggc 2760
ttggagggtg tgtttgcact gctttttgta gagagggtca cctactagaa gagaaatgcc 2820
cagctcttaga ggtgggtcag gtgtagagct ggagggggtc cctggctgct gaggggaccc 2880
taccagatga gccctgcctc tgggagcccc cttaggaagca ccagcctgga cctaccacct 2940
gcgagggcct gctgccccct ggcggccagt gctgttagag tgctgccaag cacagcctta 3000
ttcttgccgg ggccctccca ccggagagcc cagggggccg gccgggttcc tggtccttgg 3060
ctgggagcag ggctttctgg tagttggggc acaaaacat cggggaacca catgttgact 3120
gtgagcaaaag tgtcttccga ttagcagcct cagggatgcc ctggtggcct ctccagggtc 3180
gctcaggcaa ggccccccac ccattctggtg tggaaacctg ccggtccag gccagacca 3240
ggagccaaga gaaggctgaa gccagcttgg ctgtgttctc tgatctaggc cttcccagag 3300
gaggcgagca gaagctgtgc cacttggaaat tgcaacccat gagttcagaa ggcacactc 3360
gccatgctga gctccaaggg tgcctaccag ggaagatggg atctatagag tctctggggc 3420
ctggccccag ggaggagcac atttttcttg accctcacct acctggtgct agttggtcaa 3480
ccctgcctgc atacatgggc tctgtcatg gggcccagag tcccttgag atatagaat 3540
aggggaggag ctcattctgg cgccaggcag gaagaaggca ggcttctggc ttccagaggt 3600
gccgcggtgg cctcctggca tcatttgtta ttgctctga aacaagcctt actgcctgga 3660
gggcttagat tctgtcttct ccaatgtagt ttgggtatct tgtaggggat gtgggtgatg 3720
ccagggcggtg tcccaggcac ctcttctctga agtctctgca ttggagatt cgtggagaac 3780
ctatttaagc ccaattttaa ctgaaagcca gtgagctga tatggaagg aatgtaaaat 3840
ttgcttgact tcttaagaac aaaaccccca gctctgtgcc ccatgctcct tggggcttgc 3900
caccactcc tttgtgtca gaggtacagg agctgggag 3939

```

<210> 18
 <211> 1381
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6383934CB1

```

<400> 18
atgaggacaa tgctgtctgg cccacatgac ggggggatgt agacggcagc ggcgccagtc 60
gctcctggca ccatggacga tgccacagtc ctaaggaaga agggttacat cgtaggcatc 120
aatcttggca aggtttccta cgcaaaagtc aaatctgcct actctgagcg cctcaagttc 180
aatgtggctg tcaagatcat cgaccgaag aaaacaccta ctgactttgt ggagagattc 240
cttctcggg agatggacat cctggcaact gtcaaccacg gctccatcat caagacttac 300
gagatctttg agacctctga cggacggatc tacatcatca tggagcttgg cgtccagggc 360
gacctctcgc agttcatcaa gtgccaggga gccctgcatg aggacgtggc acgcaagatg 420
ttccgacagc tctctccgc cgtcaagtac tgccacgacc tggacatcgt ccaccgggac 480
ctcaagtgcg agaaccttct cctcgacaag gacttcaaca tcaagctgtc tgactttggc 540
ttctccaagc gctgcctgcg ggacagcaat gggcgcatca tctcagcaa gaccttctgc 600
gggtcggcag catatgcagc ccccgaggtg ctgcagagca tcccctacca gccaagggtg 660
tatgacatct ggagcctagg cgtgatcctc tacatcatgg totgcggtc catgccctac 720
gacgactccg acatcaagaa gatgtcgtcgt atccagaagg agcaccgcgt caacttccca 780
cgctccaagc acctgacctg cgagtgaag gacctcatct accacatgct gcagcccgac 840
gtcagccagc ggtccacat cgatgagatc ctcagccact cgtggctgca gcccccaag 900
cccaaagcca cgtcttctgc ctcttcaag agggaggggg agggcaagta ccgcgctgag 960
tgcaaaactg acaccaagac aggtctgagg cccgaccacc ggcccagca caagcttggg 1020
gcaaaaaccc agcaccggct gctggtggtg cccgagaacg agaacaggat ggaggacagg 1080
ctggccgaga cctccagggc caaagaccat cacatctccg gagctgaggt ggggaaagca 1140
agcacctagc atgacaatgg ccccgcttgtg tgtggtgggg gtccggggtg gggggcatgg 1200
tgcagtcggc cttcacgtaa actaagtagg caggtaggat ctgaagaagg cacaggtgca 1260
agtaaaatc gtcaattaaa ccactatttt gattacgttc cattagcttt cttccactta 1320
gcagcaaaag cgttccttac tgaccaccaa ataaaccaca ggggtgtgtgc aagcatcaaa 1380
a 1381

```

<210> 19
 <211> 3904
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID No: 3210906CB1

<400> 19

tattcgggggt	tcagacccca	caatcagaaa	tccggaattc	ggcagctgtc	gccctcgacg	60
aggggggagga	ctggaccgcg	aggtcagatt	aggttgtcac	cccctcccct	ccaggggagg	120
cttcccgggc	ccgcccctca	ggaagggcga	aagccgagga	agaggtggca	aggggaaagg	180
tctccttgcc	cctctccctg	acttggcaga	gccgctggag	gaccccaggc	ggaagcggag	240
gcgctggggc	accatagtg	cccctaccag	gccaggcccc	actctcaggg	ccccaggggg	300
ccaccatgcc	agctgggggc	cggggcggga	gcctgaagga	cccagatgtg	gctgagctct	360
tcttcaagga	tgacccagaa	aagctcttct	ctgacctccg	ggaaattggc	catggcagct	420
ttggagccgt	atactttgcc	cgggatgtcc	ggaatagtga	ggtggtggcc	atcaagaaga	480
tgtcctacag	tgggaagcag	tccaatgaga	aatggcaaga	catcatcaag	gaggtgcggt	540
tcttacagaa	gctccggcat	cccaacacca	ttcagtaccg	gggctgttac	ctgaggggagc	600
acacggctct	cctgttaatg	gagtattgcc	tgggetcaac	ttctgacctt	ctagaagtgc	660
acaagaaacc	gcttcaggag	gtagagatcg	cagctgtgac	ccacggggcg	cttcaggggcc	720
tggcatatct	gcactcccac	aacatgatcc	atagggatgt	gaaggctgga	aacatcctgc	780
tgtcagagcc	agggttagt	aagctagggg	actttggttc	tgctgccatc	atggcacctg	840
ccaactcctt	ctggggcacc	ccaacttgga	tggcaccoga	ggtgatccct	gccatggatg	900
aggggcagta	cgatggcaaa	gtggacgtct	ggtccttggg	gataacctgc	atcgagctgg	960
ctgaacggaa	accaccgctc	tttaacatga	atgcgatgag	tgctttatac	cacattgcac	1020
agaacgaatc	ccccgtgctc	cagtcaggac	actggtctga	gtacttccgg	aattttgtcg	1080
actcctgtct	ccagaacata	cctcaagaca	gaccaacctc	agaggttctc	ctgaagcacc	1140
gcttttgtct	ccgggagcgg	ccaccacag	tcatcatgga	cctgatccag	aggaccaagg	1200
atgccgtgct	ggagctggac	agcctgcagt	accgcaagat	gaagaagatc	ctgttccaag	1260
aggcacccaa	cggccctggt	gccgagggcc	cagaggagga	agaggaggcc	gagccctaca	1320
tgcaactctg	cgggaactct	accagcctcg	agagttagcca	ctcagtgcct	agcatgtcca	1380
tcagcgcctc	cagccagagc	agctccgtca	acagcctagc	agatgcctca	gacaacgagg	1440
aagaggagga	ggaggaggag	gaagaggagg	aggagggaaga	agggccctgaa	gccccgggaga	1500
tggccatgat	gcaggagggg	gagcacacag	tcacctctca	cagctccatt	atccaccggc	1560
tgcctgggct	tgacaacctc	tatgatgacc	cctaccagcc	agagataacc	cccagccctc	1620
tccagccgcc	tgcagcccca	gctcccactt	ccaccacctc	ttccgcccgc	cgccgggcct	1680
actgccgtaa	ccgagaccac	tttgcaccca	tccgaaccgc	ctccctggtc	agccgtcaga	1740
tccaggagca	tgagcaggac	tctgcgtgct	gggagcagct	gagcggctat	aagcggatgc	1800
gacgacagca	ccagaagcag	ctgctggccc	tggagtcacg	gctgaggggt	gaacgggagg	1860
agcacagtgc	acggctgcag	cgggagcttg	aggcgcagcg	ggctggcttt	ggggcagagg	1920
cagaaaagct	ggcccggcgg	caccaggcca	taggttgagaa	ggaggcacga	gctgcccagg	1980
ccgaggagcg	gaagtccag	cagcacatcc	ttgggcagca	gaagaaggag	ctggctgccc	2040
tgctggagct	acagaaagcg	acctacaaac	ttcgcaagga	acagctgaag	gaggagctcc	2100
aggagaaccc	cagcactccc	aagcgggaga	aggccgagtg	gctgctgcgg	cagaaggagc	2160
agctccagca	gtgccaggcg	gaggagggaag	cagggctgct	gcgggcggcag	cgccagtact	2220
ttgagctgca	gtgtcgccag	tacaagcgca	agatgttgct	ggctcggcac	agcctggacc	2280
aggacctgct	gcgggaggac	ctgaacaaga	agcagaccca	gaaggacttg	gagtgtgcac	2340
tgctgcttcg	gcagcacgag	gccacgcggg	agctggagct	gcggcagctc	caggccgtgc	2400
agcgcacgcg	ggctgagctc	accgcctgct	agcaccagac	ggagctgggc	aaccagctgg	2460
agtacaacaa	gcggcgctgag	caagagtgtc	ggcagaagca	tgcgggccag	gttcgcccag	2520
agcccaagag	cctcaaatct	aaggagctgc	agatcaagaa	gcagttccag	gagacgtgta	2580
agatccagac	tcggcagtac	aaggctctgc	gagcacactt	gctggagacc	acgcccacaa	2640
ctcagcacaa	gagcctcctt	aagcggctca	aggaagagca	gacccgcaag	ctggcgatct	2700
tggcggagca	gtatgaccag	tccatctcag	agatgtctag	ctcacaggcg	ctgcggcttg	2760
atgagaccca	ggaggcagag	ttccaggccc	ttcggcagca	gcttcaacag	gagctggagc	2820
tgctcaacgc	ttaccagagc	aagatcaaga	tccgcacaga	gagccagcac	gagaggggagc	2880
tgcgggagct	ggagcagagg	gtcgcgctgc	ggcgggcact	gctggagcag	cgggtgggaag	2940
aggagctgct	ggccctgcag	acaggacgct	ccgagcgaat	ccgcagctcg	cttgagcggc	3000
aggcccgtag	gatcgaggcc	ttcgatcgcg	aaagcatgag	gctgggcttc	tccagcatgg	3060
ctctgggggg	catcccggtc	gaagctgctg	cccaggggcta	tctgtctcca	ccccctgccc	3120
cagcctggcc	ctcccgtccc	gttccccgtt	ctggggcaca	ctggagccat	ggccctcctc	3180
caccaggcat	gccccctcca	gcctggcgct	agcgtctct	gctggctccc	ccaggccccc	3240
caaaactggct	ggggccccc	acacaaagt	ggacaccccc	tggcggagcc	ctgctgctgc	3300
taagaaacag	ccccagccc	ctgcccgggg	cagcctcggg	gggcagtgcc	agtgaagaatg	3360
tgggcccccc	tgtcgccggc	gtgcccgggc	ccctgagccg	cagcaccagt	gtcgttccc	3420
acatctcaa	tggttcttcc	cacttctatt	cctgaggtgc	agcggggagg	agcagatgag	3480
ctgggcaggg	cagggtggg	tggagcctga	ccctggaggg	actgagctg	gaggcccttg	3540
caagggtagg	ggacaagatg	taggctccag	ctcccctcag	acctcctcat	ctcatgagct	3600
tcttggggct	ggccagtggc	ccaggggccag	cttggcgata	gatgcctcaa	ggctgcctgg	3660
gagccccggc	tcccatacat	ggtgccaggg	gtctccctcc	gccacctagg	aaaggaggga	3720
gatgtcgctg	tcaaatattc	atctagctcc	ctgggggagg	ggaaaggtgg	gtctagacat	3780
actatatcca	gagaactata	ctaccctcac	agtgaggccc	tcagacctgc	cacagggcag	3840

agcaggctctg gggcctgagg cagggagaat gagaggccac ttactggcag gaaggatcag 3900
gatg 3904

<210> 20
<211> 1987
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 3339024CB1

<400> 20
gaagaaccct gaggaacaga cttacctcag caaccctggc acctccaacc cgacacatgc 60
tactgctgct gctactgctg ccaccctgc tctgtgggag agtgggggct aaggaacaga 120
aggattacct gctgacaatg tagaagtcgc tgacgggtgca ggagggcctg tgtgtctctg 180
tgctttgtct cttctcctac ccccaaaatg gctggactgc ctccgatcca gttcatggct 240
actggttccg ggcaggggac catgtaagcc ggaacattcc agtggccaca aacaaccag 300
ctcgagcagt gcaggaggag actcgggacc gattccacct ccttggggac ccacagaaca 360
aggattgtac cctgagcatc agagacacca gagagagtga tgcagggaca tacgtctttt 420
gtgtagagag aggaaatatg aaatggaatt ataaatatga ccagctctct gtgaatgtga 480
cagcgctccc ggacctactg tcaagataca ggctggagggt gccagagtcg gtgactgtgc 540
aggaggggtct gtgtgtctct gtgccctgca gtgtccttta ccccatctac aactggactg 600
cctctagccc tgtttatgga tcctgggttca aggaaggggc cgatatacca tgggatattc 660
cagtggccac aaacacccca agtggaaaag tgcaagagga taccacgggt cgattctctc 720
tccttgggga cccacagacc aacaactgct ccctgagcat cagagatgcc aggaaggggg 780
attcagggaa gtactacttc caggtggaga gaggaagcag gaaatggaac tacatatatg 840
acaagctctc tgtgcatgtg acagccctga ctcacatgcc caccttctcc atcccgggga 900
ccctggagtc tggccacccc aggaacctga cctgctctgt gccctgggac tgtgaacagg 960
ggacgcccc cagcatcacc tggatggggg cctccgtgtc ctccctggac cccactatca 1020
ctcgtctctc gatgtcagc ctcacccac agccccagga ccatggcacc agcctcacct 1080
gtcagggtgac cttgcctggg gccggcgtga ccatgaccag ggctgtccga ctcaacatat 1140
cctatcctcc tcagaacttg accatgactg tcttccaagg agatggcaca gcatccacaa 1200
ccttgaggaa cgtctcgcc ctttcagtcc tggaggggcca gtcctgcaac cttgtctgtg 1260
ctgtcgacag caatccccct gccaggctga gctggacctg ggggagcctg accctgagcc 1320
cctcacagtc ctgcaacctt ggggtgctgg agctgcctcg agtgcattgt aaggatgaag 1380
gggaattcac ctgcccagct cagaacctc taggctccca gcacatttcc ctgagcctct 1440
ccttgcaaaa cgagtacaca ggcaaaatga ggccctatc aggagtgcg ctaggggcat 1500
tcgggggagc tggagccaca gccctggtct tcctgtactt ctgcatcatc ttcgtttag 1560
tgaggtcctg caggaagaaa tcggcaaggc cagcagtggt cgtgggggat acaggcatgg 1620
aggacgcaaa cgctgtctgg ggctcagcct ctcagggacc cctgattgaa tccccggcag 1680
atgacagccc cccacacat gctccgccag ccctggccac cccctccca gaggaaggag 1740
agatccagta tgcattcctc agcttcacaa aagcgaggcc tcagtacca caggaaacagg 1800
aggccatcgg ctatgagtac tccgagatca acatccccaa gtgagaaact gcagagactc 1860
aggcctgttt gagggctcac gaccctccca gcaaagaagc ccgagactga ttcctttaga 1920
attaaaagcc ctccatgctg tgcaacgggg gatccactag ttaagagcgg cgaccccgcg 1980
tgccccct 1987

<210> 21
<211> 3925
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4436929CB1

<400> 21
ccgtcctcga ggcgaggaga gtaccggggc ggcccggctg ccgcgcgagg agcgcggtcg 60
gcggcctggt ctgcggtgta gatacacaga gcgacagaga catttattgt tatttgtttt 120
ttggtggcaa aaagggaaaa tggcgaacga ctcccctgca aaaagtctgg tggacatcga 180
cctctcctcc ctgcgggatc ctgctgggat ttttgagctg gtggaagtgg ttggaaatgg 240
cacctatgga caagtctata agggctcgaca tgttaaaacg ggtcagttgg cagccatcaa 300
agttatggat gtcactgagg atgaagagga agaaatcaaa ctggagataa atatgctaaa 360
gaaatactct catcacagaa acattgcaac atattatggt gctttcatca aaaagagccc 420
tccaggacat gatgaccaac tctggcttgt tatggagtgc tgtggggctg ggtccattac 480
agacctgtg aagaacacca aagggaacac actcaaagaa gactggatcg cttacatctc 540

```

cagagaaatc ctgaggggac tggcacatct tcacattcat catgtgattc accgggatat 600
caagggccag aatgtgttgc tgactgagaa tgcagaggtg aaacttggtg acttttggtg 660
gagtgtctag ctggacagga ctgtggggcg gagaaatacg ttcataaggca ctccctactg 720
gatggctcct gaggtcatcg cctgtgatga gaaccagat gccacctatg attacagaag 780
tgatctttgg tcttgtggca ttacagccat tgagatggca gaaggtgctc cccctctctg 840
tgacatgcat ccaatgagag cactgtttct cattcccaga aaccctcctc cccggctgaa 900
gtcaaaaaaa tggctgaaga agtttttttag ttttatagaa ggttgcttgg tgaagaatta 960
catgcagcgg ccctctacag agcagctttt gaaacatcct tttataaggg atcagccaaa 1020
tgaaggcga gttagaatcc agcttaagga tcatatagat cgtaccagga agaagagagg 1080
cgagaaagat gaaactgagt atgagtacag tgggagttag gaagaagagg aggaagtggc 1140
tgaacaggaa ggagagccaa gttccattgt gaacgtgcct ggtgagtcta ctcttcgccc 1200
agatttcctg agactgcagc agggagaacaa ggaacgttcc gaggctcttc ggagacaaca 1260
gttactacag gagcaacagc tccgggagca ggaagaatat aaaaggcaac tgctggcaga 1320
gagacagaag cggagttagc agcagaaaga acagaggcga cggctagaag agcaacaaag 1380
ggagagcgg ggagctagaa ggcagcagga acgtgaacag cgaaggagag aacaagaaga 1440
aaagaggcgt ctagaggagt tggagagaag gcgcaaagaa gaaggaggaga ggagacgggc 1500
agaagaagaa aagaggagag ttgaagaga acaggagtat atcaggcgac agctagaaga 1560
ggagcagcgg cacttggaag tccttcagca gcagctgctc caggagcagg ccatgttact 1620
gcatgaccat aggaggccgc acccgcagca ctgcgcagc cgcgccaccac cgcagcagga 1680
aaggagcaag ccaagcttcc atgtctccga gcccaaagcc cactacgagc ctgctgaccg 1740
agcgcgagag gtggaagata gatttaggaa aactaaccac agctcccctg aagcccagtc 1800
taagcagaca ggcagagtat tggagccacc agtgccctcc cgatcagagt ctttttccaa 1860
tgccaactcc gagtctgtgc atccgcctc gcagagacca gcggagccac aggttccctg 1920
gagaacaaca tctcgtctcc ctgttctgtc ccgtcgagat tcccactgc agggcagtg 1980
gcagcagaat agccaggcag gacagagaaa ctccaccagc agtattgagc ccaggcttct 2040
gtgggagaga tgggagaagc tgggtcccag acctggcagt ggcagctcct caggggtccag 2100
caactcagga tcccagcccg ggtctcacc tgggtctcag agtggctccg gggaaacgct 2160
cagagtgaga tcatcatcca agtctgaagg ctctccatct cagcgcctgg aaaatgcagt 2220
gaaaaaacct gaagataaaa aggaagtttt cagaccctc aagcctgctg gcgaagtgga 2280
tctgaccgca ctggccaaag agcttcgagc agtggaagat gtacggccac ctcaaaagt 2340
aacggactac tcctcatcca gtgaggagtc ggggacgacg gatgaggagg acgacgatgt 2400
ggagcaggaa ggggctgacg agtccacctc aggaccagag gacaccagag cagcgtcatc 2460
tctgaatttg agcaatggtg aaacggaatc tgtgaaaacc atgattgtcc atgatgatgt 2520
agaaagtggc ccggccaatg agtctcacc ccccatccaa ctaatcgtcc gccagactca 2580
gtccgctagt agcacactcc agaaacacaa atcttctctc tcctttacac cttttataga 2640
ccccagatta ctacagattt ctccatctag cggaaacaaca gtgacatctg tgggtgggatt 2700
ttcctgtgat gggatgagac cagaagccat aaggcaagat cctaccgga aaggctcagt 2760
ggtaaatgtg aatcctacca acactaggcc acagagttag accccggaga ttcgtaata 2820
caagaagagg tttaactctg agattctgtg tgctgcctta tggggagtga atttgctagt 2880
gggtacagag agtggcctga tgctgctgga cagaagtggc caaggggaagg tctatcctct 2940
tatcaaccga agacgatttc aacaaatgga cgtacttgag ggcttgaaatg tcttgggtgac 3000
aatatctggc aaaaaggata agttacgtgt ctactatttg tcctggttaa gaaataaaat 3060
acttcacaa gatccagaag ttgagaagaa gcagggatgg acaaccgtag gggatttgga 3120
aggatgtgta cattataaag ttgtaaaata tgaaagaatc aaatttctgg tgattgcttt 3180
gaagagtctt gtggaagtct atgctggggc accaaagcca tatcaciaat ttatggcctt 3240
taagtcatct ggagaattgg attactgggt tgctggattc catgctgttg atgtggattc 3300
ccagaggttg aaagtgatct atggatcctg tgctggattc catgctgttg atgtggattc 3360
aggatcagtc tatgacattt atctaccaac acatatccag tgtagcatca aaccccatgc 3420
aatcatcatc ctccccaaata cagatggaat ggagcttctg gtgtgctatg aagatgaggg 3480
ggtttatgta aacacatatg gaaggatcac caaggatgta gttctacagt ggggagagat 3540
gcctacatca gtagcatata ttcgatccaa tcagacaatg ggctggggag agaaggccat 3600
agagatccga tctgtggaaa ctgggtcactt ggatgggtgtg ttcatgcaca aaagggctca 3660
aagactaaaa ttctgtgtg aacgcaatga caagtggttc tttgcctctg ttcggtctgg 3720
tggcagcagt caggtttatt tcatgacctt aggcaggact tctcttctga gctggtagaa 3780
gcagtgtgat ccagggatta ctggcctcca gagtcttcaa gatcctgaga acttggaatt 3840
ccttgtaact ggagctcgga gctgcaccga gggcaaccag gacagctgtg tgtgcagacc 3900
tcatgtgttg ggttctctcc cctcc

```

<210> 22

<211> 1210

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5046791CB1

WO 01/46397

<400> 22
ttacagggtca tctaccctta taccacacaa aatgacgatg
gacttcatct tcatgtctcc aatggagcag accagcacca gcy
acgtccctaa ccaccggctg ctctggactc ctcttgagaa ttaca
aatgcagcac ctggatattt catgttctt attcaatctt tgaggacc
ctctcacgtt tggggatgga gtattggaga ggcggcctta gctgagggtc
agacgactcc tcttactatc atctggcagc ccatgcagcc tgagaggatg
ccggcccccga gaagcgatgc cagtgtctcg atgcccagg cgcctacata cgca
ttgggaagta ctggctgtcc cgtgtgctcg atgcccagg cgcctacata cgca
tgaacatgcc tcatagtcta cctcagcaga gtgggtggtt cgcgagattc gagaaa
ctcccatcac tgtgtttgga tgcattgct cccgtccct ggcgtgctt cagactgca
gcaataacct gaaaggttta caacaagaaa atcacttgaa gatccgtgta gagcccgt
tatttagtg gacaaaatgg gttgctggga gcacattacc tgcatggata cctccatcag 780
agttagctgc agccaaactg agtgttgata caacctacag acctcatt ccaatcagca 840
aattagtgt ttcagaaatc tatgatact atactagtag aagtttcaa gtaacaaaag 900
aaataataag tgaatgtaaa agtaaaaggaa ataactcct gattgtggcc cagcatctt 960
cccttgaagc gtgtacctgc caacttcagg gctgtcacc tcagaactcc aaggacttcg 1020
tacaataggt ccgaaagatc ccatactgg gattttgtc ctgtgaagaa ttaggagaaa 1080
ctggaaatag gcagctgaca gatccacaa tcttctctt taccatgga ccaactggg 1140
gttcaactg gagagagacc ttgcttcaag aataaaccat accagtgaac aagaaggaaa 1200
aaaaaaaaa 1210

<210> 23
<211> 1521
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1416174CB1

<400> 23
ggcagcgtgc tgggcctcgc cacactgcac acctaccgag gacgcctctc ctacctccc 60
gccactgtgg aacctgcctc gccaccct gccatagcc tgccctgtgc caagtcggag 120
ctgacctaa cccagaccc agcccgccc atggccact caccctgca tctgtctgtg 180
tctgacctgc ctctccctc cgggtggggc ccagagctgg ctggggactg ggggtggggc 240
ctgccccatc tgtccctcaa ggaacactg ctgtcttcac ctctggctc tcccaaggca 300
ggggatgctc cgtgtccc cgaaggccc ccactgcgg ccgcccagc cactgtctgc 360
gctctacact caccgtctc gtagggcct ccactgcgg ccgcccagc cactgtctgc 420
ttccacccc ttagtccc cccagagct ggtgacgct ggcagctcc catgctgctg 480
ctccgctggg caccgctgc cactaggg ctgacctgg tgcgtacgg catctcggc cgcagctg 540
tggtggcat ctgcggcag cctgtgtgg tgcgtacgg catctcggc cgcagctg 600
tgcgctttt ctggccatg gagcgtgga gccacttcag cctgggctg cgcagctg 660
gctacgccc ggcctgtgc tatggccc ggcgggagc ctgaaactaa acaagcttg 720
tggacgggga gcaggtggg gctgcccgg tggggcgag ctgagctagg ggtgtgtgg 780
cactgtcac tgggctcct tacattcaa tggcccctg tggcccctg aatgggctg 840
taccgcccgg gggcggggc gcaggggccc cgtcgtcac gatgaccct ggcagctg 900
tggtgctag agttgtgtg ggaaggtgg cgtcgtcac gatgaccct ggcagctg 960
atgggaccag acgtgatgt caatgaggg gggccctgg taccggcca ggtgtgtg 1020
tccgaggggt agtcctgat tgacgcttc gacaggacct ggaatgtact ggtgtgtg 1080
tacgggggagc tattttacg tgcggccc cagcctcgc ccatccactc cgtgtgtg 1140
ggcctcagtg agtcggccg tcaggggccc aggttcccc gggcgggcg taggtgtg 1200
atttagctg ccaatcagc caggagggg ggggaaattc atatccctg ttcgtctc 1260
actaatgtc ctctccccg ggggtggggc attgaaaagg tctatgcaat aaaggcagtc 1320
gcgctctc cgtcccaat ctaaaaagca attgaaaagg tctatgcaat aaaggcagtc 1380
gcttcattc tctcaaaaa a 1440
1500
1521

<210> 24
<211> 1640
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature

WO 01/46397

<223> Incyte ID No: 3244919CB1

0/35304

<400> 24
gcagcgccgc ggcgtccccg ggctcgccgc ccccgccgc cgcgcgcccc g
acgcgcctc ggcctcgccg ccgcccgtg ctggccagcc ccgggcccgg gac
atgtccgctc gcagccgcgc cccctgttcc agtggagcaa gtggaagaag agga
cgtccatgtc ggcggccacc gcgcggaggc cgtgtttga cgacaaggag gacgtg
tcgaccactt ccagatcctt cgggccattg ggaaggagc ctttgcaag gtgtgca
tcgagaagcg ggacacggag aagatgtacg ccatgaagta catgaacaag cagcagtgc
tcgagcgcgga cgaggtcgc aacgtcttcc tccaggacga ggaggacatg gagatcgagc
acgtcttct ggtgaacctc tggactctc accacctgca gcagaacgtg cagttctccg 46
tggacctgct actgggcccgc gacctgcgt tggactggc tctggactac ctgcgcggcc 540
aggacacggt gaggctgtac atctgcgaga tggactggc cctggatgag agaggacatg 600
agcacatcat ccacagagat gtcaagcctg acaacattct cctggatgag agaggacatg 660
cacacctgac cgacttcaac attgccacca tcatcaagga cggggagcgg gcgacggcat 720
tagcaggcac caagccgtac atggctccg agatcttcca ctctttgtc aacggcggga 780
ccggctactc cttcgaggtg gactggtggt cgggtggggg gatggcctat gtcagctgt 900
tcagaccgt gaggctccag tatgtccca gcaacggcgt ggagtccctg gccttgctgc 960
ggaagctct cactgtgaac ccgagcacc ggctctccag cctccaggac gtgcaggcag 1020
ccccggcgt ggcggcgtg ctgtgggacc accccacctt tgagctggag gagecgggct 1080
tcgtgccca caaaggcgt ctgcactgc agccctggc caagaacaag tcccgggaca 1140
tggagtccag gcccctgcac aagaagaaga agccctggc caagccatcc agactgcctc 1200
acagcaggga cagctcccag tccgcccac ggagcaagtc actatcttca agaccaggac 1260
aaggagctg ggccttgga tcctcgggct tgggagaatg agctgaagag gaggcaggac 1320
gatgccatcc agcaagactt cgtgattttt aacagagaaa agctgaagag tgtggaggac 1380
ctcccgagg agcctctccc cgcctctgag tccagggatg ctgcggagcc tgtggaggac 1440
taggcccgaac gtcctgcctt gccatgtgc ggccccattt gcccctcggc cgggagcggc 1500
ggaggggccat ggcccgaggc ctggcattca cgttccacc cagcctggct ggcggtgccc 1560
acagtgcctt ggacacattt 1620
1640